

12-30-2020

Effect of Cultivation Parameters on Fermentation and Hydrogen Production in the Phylum

Mariamichela Lanzilli
Istituto di Chimica Biomolecolare

Nunzia Esercizio
Istituto di Chimica Biomolecolare

Marco Vastano
Istituto di Chimica Biomolecolare

Zhaohui Xu
Bowling Green State University

Genoveffa Nuzzo
Istituto di Chimica Biomolecolare

Follow this and additional works at: https://scholarworks.bgsu.edu/bio_sci_pub

See next page for additional authors



Part of the [Biology Commons](#)

Repository Citation

Lanzilli, Mariamichela; Esercizio, Nunzia; Vastano, Marco; Xu, Zhaohui; Nuzzo, Genoveffa; Gallo, Carmela; Manzo, Emiliano; Fontana, Angelo; and d'Ippolito, Giuliana, "Effect of Cultivation Parameters on Fermentation and Hydrogen Production in the Phylum" (2020). *Biological Sciences Faculty Publications*. 92.

https://scholarworks.bgsu.edu/bio_sci_pub/92



This work is licensed under a [Creative Commons Attribution 4.0 International License](#).

This Article is brought to you for free and open access by the Biological Sciences at ScholarWorks@BGSU. It has been accepted for inclusion in Biological Sciences Faculty Publications by an authorized administrator of ScholarWorks@BGSU.

Author(s)

Mariamichela Lanzilli, Nunzia Esercizio, Marco Vastano, Zhaohui Xu, Genoveffa Nuzzo, Carmela Gallo, Emiliano Manzo, Angelo Fontana, and Giuliana d'Ippolito



Review

Effect of Cultivation Parameters on Fermentation and Hydrogen Production in the Phylum *Thermotogae*

Mariamichela Lanzilli ¹, Nunzia Esercizio ¹, Marco Vastano ¹, Zhaohui Xu ², Genoveffa Nuzzo ¹, Carmela Gallo ¹, Emiliano Manzo ¹, Angelo Fontana ¹ and Giuliana d'Ippolito ^{1,*}

¹ Istituto di Chimica Biomolecolare (ICB), CNR, Via Campi Flegrei 34, 80078 Pozzuoli, Napoli, Italy; mariamichelanzilli@gmail.com (M.L.); esercizionunzia@gmail.com (N.E.); marco.vastano@gmail.com (M.V.); nuzzo.genoveffa@icb.cnr.it (G.N.); carmen.gallo@icb.cnr.it (C.G.); emanzo@icb.cnr.it (E.M.); afontana@icb.cnr.it (A.F.)

² Department of Biological Sciences, Bowling Green State University, Bowling Green, OH 43403, USA; zxu@bgsu.edu

* Correspondence: gdippolito@icb.cnr.it; Tel.: +39-081-8675096

Abstract: The phylum *Thermotogae* is composed of a single class (*Thermotogae*), 4 orders (*Thermotogales*, *Kosmotogales*, *Petrotogales*, *Mesoaciditogales*), 5 families (*Thermatogaceae*, *Fervidobacteriaceae*, *Kosmotogaceae*, *Petrotogaceae*, *Mesoaciditogaceae*), and 13 genera. They have been isolated from extremely hot environments whose characteristics are reflected in the metabolic and phenotypic properties of the *Thermotogae* species. The metabolic versatility of *Thermotogae* members leads to a pool of high value-added products with application potentials in many industry fields. The low risk of contamination associated with their extreme culture conditions has made most species of the phylum attractive candidates in biotechnological processes. Almost all members of the phylum, especially those in the order *Thermotogales*, can produce bio-hydrogen from a variety of simple and complex sugars with yields close to the theoretical Thauer limit of 4 mol H₂/mol consumed glucose. Acetate, lactate, and L-alanine are the major organic end products. *Thermotogae* fermentation processes are influenced by various factors, such as hydrogen partial pressure, agitation, gas sparging, culture/headspace ratio, inoculum, pH, temperature, nitrogen sources, sulfur sources, inorganic compounds, metal ions, etc. Optimization of these parameters will help to fully unleash the biotechnological potentials of *Thermotogae* and promote their applications in industry. This article gives an overview of how these operational parameters could impact *Thermotogae* fermentation in terms of sugar consumption, hydrogen yields, and organic acids production.

Keywords: anaerobic bacteria; hydrogen yields; fermentation rate; organic acids; nitrogen; carbon dioxide



Citation: Lanzilli, M.; Esercizio, N.; Vastano, M.; Xu, Z.; Nuzzo, G.; Gallo, C.; Manzo, E.; Fontana, A.; d'Ippolito, G. Effect of Cultivation Parameters on Fermentation and Hydrogen Production in the Phylum *Thermotogae*. *Int. J. Mol. Sci.* **2021**, *22*, 341. <https://doi.org/10.3390/ijms22010341>

Received: 12 October 2020

Accepted: 23 December 2020

Published: 30 December 2020

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

The phylum *Thermotogae* is comprised of thermophilic, hyperthermophilic, mesophilic, and thermo-acidophilic anaerobic bacteria that originated from geothermally heated environments (Table 1) [1,2]. Recent phylogenetic analyses based on gene markers/core genome inferences, comparative genomics, and whole-genome relatedness have led to a taxonomic revision of the phylum, with a single class (*Thermotogae*), 4 orders (*Thermotogales*, *Kosmotogales*, *Petrotogales*, *Mesoaciditogales*), 5 families (*Thermatogaceae*, *Fervidobacteriaceae*, *Kosmotogaceae*, *Petrotogaceae*, *Mesoaciditogaceae*), and 13 genera, i.e., *Thermotoga* (T.) [3], *Pseudothermotoga* (Pseudot.) [2,4], *Fervidobacterium* (F.) [5], *Thermosiphon* (Ts.) [6], *Kosmotoga* (K.) [7], *Mesotoga* (Ms.) [8], *DeFluviitoga* (D.) [9], *Geotoga* (G.) and *Petrotoga* (P.) [10], *Marinitoga* (Mn.) [11], *Oceanotoga* (O.) [12], *Mesoaciditoga* (M.) [13], and *Athalassatoga* (A.) (Table 1) [2,4,14]. *Thermotogae* are able to grow under mesophilic (*Kosmotogales*; *Mesoaciditogales*, *Petrotogales*) and thermophilic conditions (*Thermotogales*), but most species have optimal growth temperatures in the range of 45–80 °C (Table 1). They are Gram-negative

bacteria, except for *D. tunisiensis*, which shows a positive result in Gram staining [9]. Apart from *K. shengliensis*, whose cells are in a coccoid form, *Thermotogae* cells are rod-shaped and encapsulated by a unique outer membrane, named “toga” [1,8,15]. Usually, the cells grow singly or in pairs, but it is also possible to observe chains surrounded by a unique toga [1,2]. Cell length is typically less than 20 μm , except for *F. gondwanense* and some members of the *Petrotoga* genus, whose cells can reach to 50 μm long (Table 1) [2,10]. Almost all species grow at neutral pH, and NaCl tolerances are high among *Geotoga*, *Oceanotoga*, and *Petrotoga* species (Table 1). Numerous studies have reported that members of the phylum can grow on both simple (e.g., glucose, galactose, fructose, lactose, maltose, mannose, sucrose) and complex carbohydrates (e.g., starch, glycogen, cellulose, keratin) (Table 1). Genes, transcriptional factors, and regulatory mechanisms driving the carbohydrates utilization have been identified for multiple members of the phylum [16–18]. ABC transporters for the uptake of a broad list of sugars have also been characterized [19–23].

All species of the phylum, except for *Mesotoga spp.*, have tremendous potentials in biotechnological production of H_2 , especially the order *Thermotogales*, as their hydrogen yields are close to the theoretical maximum value (Thauer limit) of 4 mol H_2 /mol glucose [1,4,24]. Acetate, lactate, and L-alanine are the major organic products of the sugar fermentation [1]. *Ms. prima* and *Ms. infera* produce mainly/only acetate from sugar utilization without H_2 formation [8,25–27]. Lactate is produced by *T. maritima*, *T. neapolitana*, and *Mn. camini* in variable quantities depending on growth conditions [11,28–31]. Other significant products include ethanol (has been measured in *Geotoga*, *Petrotoga*, *Kosmotoga*, and *Oceanotoga spp.*); isovalerate, isobutyrate, and/or propionate (have been measured in *Mn. camini* and *K. olearia*); L-glutamate, alpha-aminobutyrate, hydroxyphenyl-acetate, or phenylacetate (have been measured in *F. pennavorans*) [1,32] (Table 1). Among these fermentation end-products, lactic acid has been widely used in various industries such as food, cosmetic, pharmaceutical, and chemical industries, although its primary application is serving as the building block for the production of biodegradable polylactic acid (PLA) [33]. Ethanol is an important industrial commodity; it is used as a food additive and a renewable biofuel; it is also contained in many cosmetics, households, and sanitizer products [34]. Moreover, a plethora of thermostable enzymes, harbored by most of these bacteria, are valuable components for many industrial and biotechnological applications [17,35–44].

Hydrogen (H_2) is considered a green and sustainable alternative to traditional fossil fuels and is capable of mitigating greenhouse gas emissions. Using hydrogen in fuel cells or combustion engines produces heat and electricity with water as the only waste. As the current abiotic hydrogen production method is energy-consuming and still causes pollution, emphasis must be given to biological production of the energy from renewable sources [45,46]. Biological synthesis of H_2 can use a wide range of organic substrates as feedstocks, including agro-industrial wastes and algal biomass, and may operate under various environmental conditions [1,46–54]. In addition, high temperatures help to improve the solubilization of substrates, reduce fermentation time, and lower contamination risks [55]. Although hydrogen production by *Thermotoga* species is considered one of the most challenging biological systems, no application using pure *Thermotoga* cultures has been reported at the industrial scale.

Releasing hydrogen is an efficient way to dissipate excessive reductants generated during the fermentative conversion of organic substrates. The process is generally referred to as dark fermentation (DF) and is typically influenced by environmental conditions such as pH, cell growth rate, and hydrogen partial pressure [24,56,57].

According to the classical model of dark fermentation, theoretically up to 4 mol of hydrogen may be produced from each mole of glucose, which is converted to acetate and CO_2 (Thauer limit Figure 1) [24]. When hydrogen accumulates, pyruvate is diverted away from acetate production. In this case, excessive NADH from glycolysis is not used in the energetically favorable manner to synthesize acetate and H_2 but dissipated via synthesizing other metabolic products such as lactic acid, L-alanine, ethanol, butyrate, and valerate (Figure 1) [24]. Synthesis of hydrogen in *Thermotogae* species is performed

by the heterotrimeric [FeFe]-hydrogenase, an electron-bifurcating enzyme that couples the endergonic reduction of H^+ to hydrogen by NADH to the exergonic reduction of H^+ to hydrogen by reduced ferredoxin (Figure 1) [58]. Because the hydrogenase uses both NADH and reduced ferredoxin as electron donors, hydrogen yield is influenced by factors that affect both reductants.

The value of these bacteria in biotechnological processes is rising sharply since the discovery of the bifurcating hydrogenase and will probably be enhanced with a full elucidation of the molecular and biochemical properties of the processes. Despite decades of efforts in the development of genetic tools to engineer these species, only a few of thermostable selectable markers and genetic modifications with low stability are reported, which makes it still difficult to perform genetic modifications of these organisms [59–61]. However, these difficulties could be offset by their well-known susceptibility to mutations under environmental pressures [62,63].

In recent years, many researchers have been focusing on the optimization of fermentation performance towards the production of hydrogen and other target end-products [30,43,64–71].

Anaerobic fermentation in *Thermotogae* depends on many cultivation parameters such as hydrogen partial pressure, agitation, gas sparging, culture/headspace ratio, inoculum, pH, temperature, nitrogen sources, sulfur sources, inorganic compounds, and metal ions. The effect of each factor on H_2 yield, sugar consumption rate, and formation of biotechnologically interesting end-products are discussed here. Main data are also summarized in extensive tables, citing the most important studies, with the information on their cultivation systems (e.g., reactor type, incubation periods, batch vs. continuous modality).

Table 1. Physiological and metabolic properties of *Thermotoga* species. **YE:** Yeast extract; **BHI:** Brain heart infusion; **CMC:** Carboxymethylcellulose; **S⁰** = Elemental sulfur; **Thio:** Thiosulfate; **Cys:** Cysteine; **AA:** Acetic acid; **LA:** Lactic acid; **ALA:** Alanine; **EPS:** Exopolysaccharide; **AABA:** α -aminobutyrate; **EtOH:** Ethanol; **AQDS:** Anthraquinone-2,6-disulfonate; **But:** Butyrate; **Val:** Valerate; **Glu:** Glutamate; **BuOH:** Butanol; **iBut:** isobutyrate; **iVal:** isovalerate; **PPA:** Propionic Acid; **Gly:** Glycine; **Pro:** Proline; **Fo:** Formate; **HPA:** Hydroxyphenilacetate; **PA:** Phenylacetate; **3-IAA:** Indole-3-acetate; **2-MeBu:** 2-Methylbutyrate.

Genus	Species	Isolation	Temp. Range/ Optimal (°C)	pH Range/ Optimal	Cell Dimension (Long by Wide) (μ m)	Growth Substrates	NaCl Range/ Optimal (%)	Electron Acceptor	End Products	Ref.
<i>Thermotoga</i>	<i>Thermotoga petrophila</i>	Oil reservoir, Japan	47–88/ 80	5.2–9.0/ 7.0	2.0–7.0 by 0.7–1.0	YE, peptone, glucose, fructose, ribose, arabinose, sucrose, lactose, maltose, starch, cellulose	0.1–5.5/ 1.0	S ⁰ ; Thio	AA, LA, CO ₂ , H ₂	[72]
	<i>Thermotoga naphthophila</i>	Oil reservoir, Japan	48–86/ 80	5.4–9.0/ 7.0	2.0–7.0 by 0.8–1.2	YE, peptone, glucose, galactose, fructose, mannitol, ribose, arabinose, sucrose, lactose, maltose, starch	0.1–6.0/ 1.0	S ⁰ ; Thio	AA, LA, CO ₂ , H ₂	[72]
	<i>Thermotoga maritima</i>	Geothermal vent	55–90/ 80	5.5–9.0/ 6.5	1.5–11.0 by 0.6	ribose, xylose, glucose, sucrose, maltose, lactose, galactose, starch, glycogen	0.2–3.8/ 2.7	Fe (III) S ⁰ ; Thio	AA, LA, CO ₂ , H ₂ , ALA, EPS, AABA	[3]
	<i>Thermotoga profunda</i>	Hot spring, Japan	50–72/ 60	6.0–8.6/ 7.4	0.8–2.1 by 0.4	glucose, trehalose, cellobiose, arabinose, xylose, ribose, pyruvate	n. d	S ⁰ ; Thio	n. d	[73]
	<i>Thermotoga caldifontis</i>	Hot spring, Japan	55–85/ 70	6.0–8.6/ 7.4	1.2–3.5 by 0.5	glucose, maltose, trehalose, cellobiose, arabinose, xylose, ribose, pyruvate, starch	n. d	Thio	n. d	[73]
	<i>Thermotoga neapolitana</i>	Submarine thermal vent	55–95/ 77	6.0–9.0/ 7.5	1.5–11.0 by 0.6	fructose, fucose, galactose, mannose, rhamnose, pyruvate, glucosamine, lactulose, turanose, glycerol, dextrin, ribose, xylose, glucose, sucrose, maltose, lactose, starch, glycogen	0.2–6.0/ 2.0	S ⁰	AA, ALA, CO ₂ , H ₂	[74]
<i>Pseudothermotoga</i>	<i>Pseudothermotoga lettingae</i>	Thermophilic bioreactor	50–75/ 65	6.0–8.5/ 7.0	2.0–3.0 by 0.5–1.0	glucose, EtOH, acetate, formate	0.0–2.8/ 1.0	S ⁰ ; Thio; AQDS; Fe(III)	AA, ALA, LA, EtOH, AA, BA, CO ₂ , H ₂	[75]
	<i>Pseudothermotoga elfii</i>	Oil reservoir	50–72/ 66	5.5–7.5/ 7.5	2.0–3.0 by 0.5–1.0	glucose, arabinose, fructose, lactose, maltose, mannose, ribose, sucrose, xylose	0.0–2.8/ 1.0	Thio	AA, CO ₂ , H ₂	[76]
	<i>Pseudothermotoga hypogea</i>	Oil reservoir, Africa	56–90/ 70	6.1–9.1/ 7.3–7.4	2.0–3.0 by 0.5–1.0	fructose, galactose, glucose, lactose, maltose, mannose, sucrose, xylose, xylan	0.0–1.5/ 0.2	Thio	AA, ALA, CO ₂ , H ₂ , EtOH	[77]
<i>Pseudothermotoga</i>	<i>Pseudothermotoga subterranea</i>	Oil reservoir, Paris	50–75/ 70	6.0–8.5/ 7.0	3.0–10.0 by 0.5	YE, peptone, tryptone, casein	0.0–2.4/ 1.2	Cys, Thio	n.d.	[78]
	<i>Pseudothermotoga thermarum</i>	Hot spring, Africa	55–84/ 70	6.0–9.0/ 7.0	1.5–11.0 by 0.6	starch, glucose, maltose	0.2–0.5/ 0.35	S ⁰	n.d.	[6]

Table 1. Cont.

Genus	Species	Isolation	Temp. Range/ Optimal (°C)	pH Range/ Optimal	Cell Dimension (Long by Wide) (µm)	Growth Substrates	NaCl Range/ Optimal (%)	Electron Acceptor	End Products	Ref.
<i>Ferroidobacterium</i>	<i>Ferroidobacterium nodosum</i>	Hot spring, New Zealand	40–80/ 65–70	6.0–8.0/ 7.0	1.0–2.5 by 0.5–0.55	glucose, sucrose, starch and lactose	n.d./<1.0	S ⁰	AA, LA, CO ₂ , H ₂ , EtOH, But, Val	[5]
	<i>Ferroidobacterium pennavorans</i>	Hot spring, Portugal	50–80/ 70	5.5–8.0/ 6.5	2.0–20.0 by 0.5	cellobiose, starch, glycogen, pullulan, glucose, fructose, maltose, xylose, native feathers	0.0–4.0/ 0.4	S ⁰ ; Thio	AA, CO ₂ , ALA, Glu, EtOH, But, H ₂ , BuOH	[79]
	<i>Ferroidobacterium islandicum</i>	Icelandic Hot spring	50–80/ 65	6.0–8.0/ 7.2	1.0–4.0 by 0.6	pyruvate, ribose, glucose, maltose, raffinose, starch, cellulose	0.0–1.0/ 0.2	S ⁰ ; Thio	LA, AA, H ₂ , EtOH, CO ₂ , iBut, iVal	[80]
	<i>Ferroidobacterium riparium</i>	Hot spring, Russia	46–80/ 65	5.7–7.9/ 7.8	1.0–3.0 by 0.4–0.5	peptone, YE, pyruvate, glucose, xylose, fructose, maltose, sucrose, cellobiose, starch, xylan, CMC, cellulose, filter paper	0.0–1.0/ 0.0	S ⁰	H ₂ , AA, CO ₂ , PPA, iBut, But	[81]
	<i>Ferroidobacterium gondwanense</i>	Hot spring, Australia	45–80/ 65–68	5.5–8.5/ 7.0	4.0–40.0 by 0.5–0.6	cellobiose, amylopectin, maltose, starch, dextrin, xylose, glucose, pyruvate, lactose, fructose, mannose, CMC, galactose	0.0–0.6/ 0.1	S ⁰	EtOH, AA, LA, CO ₂ , H ₂	[82]
	<i>Ferroidobacterium thailandese</i>	Hot spring, Thailand	60–88/ 78–80	6.5–8.5/ 7.5	1.1–2.5 by 0.5–0.6	glucose, maltose, sucrose, fructose, cellobiose, CMC, cellulose, starch	<0.5/0.5	S ⁰	n.d.	[83]
	<i>Ferroidobacterium changbaicum</i>	Hot spring, China	55–90/ 75–80	6.3–8.5/ 7.5	1.0–8.0 by 0.5–0.6	glucose, lactose, fructose, sucrose, maltose, starch, sorbitol, cellobiose, trehalose, galactose, melibiose, pyruvate, glycerin	0.0–1.0/ 0.0	S ⁰	n.d.	[84]
<i>Thermosipho</i>	<i>Thermosipho africanus</i>	Hot spring, Africa	53–77/ 75	6.0–8.0/ 7.2	3.0–4.0 by 0.5	glucose, ribose, maltose, starch, galactose, fructose, sucrose	0.11–3.6	S ⁰ ; Thio	AA, H ₂ , CO ₂ , LA	[85]
	<i>Thermosipho japonicus</i>	Hydrothermal vent, Japan	45–80/ 72	5.3–9.3/ 7.2–7.6	3.0–4.0 by 0.5	YE, peptone, and tryptone, maltose, glucose, galactose, starch, sacharose, ribose, casein	0.7–7.9/ 4.0	S ⁰ ; Thio	n.d.	[86]
	<i>Thermosipho geolei</i>	Oil reservoir, Russia	45–75/ 70	6.0–9.4/ 7.5	2.0–3.0 by 0.4–0.6	Glucose, peptone, beef extract, YE	0.5–7.0/ 2.0–3.0	S ⁰	H ₂ , AA, ALA, CO ₂ , iVal	[87]
<i>Thermosipho</i>	<i>Thermosipho affectus</i>	Hydrothermal vent, Atlantic Ocean	37–75/ 70	5.6–8.2/ 6.6	1.2–6.0 by 0.4–0.9	YE, beef extract, glucose, maltose, sucrose, starch, dextrin, CMC, cellulose	1.0–5.5/ 2.0	S ⁰	AA, H ₂ , CO ₂ , EtOH	[88]
	<i>Thermosipho globiformans</i>	Hydrothermal vent	40–75/ 68	5.0–8.2/ 6.8	2.0–4.0 by 0.5	YE, tryptone, starch	0.2–5.2/ 2.5	S ⁰ ; Fe ₂ O ₃	n.d.	[89]

Table 1. Cont.

Genus	Species	Isolation	Temp. Range/ Optimal (°C)	pH Range/ Optimal	Cell Dimension (Long by Wide) (µm)	Growth Substrates	NaCl Range/ Optimal (%)	Electron Acceptor	End Products	Ref.
	<i>Thermosipho melanesiensis</i>	Hydrothermal vent, Pacific Ocean	50–75/ 70	4.5–8.5/ 6.5–7.5	1.0–3.5 by 0.4–0.6	BHI, malt extract, tryptone, sucrose, starch, glucose, maltose, lactose, cellobiose, galactose	1.0–6.0/ 3.0	S ⁰	H ₂ , AA, ALA, CO ₂	[90]
	<i>Thermosipho activus</i>	Riftia sheath, Guaymas Basin	44–75/ 65	5.5–8.0/ 6.0	1.5–10.0 by 0.3–0.8	glucose, maltose, cellobiose, cellulose, filter paper, chitin, xylan, pectin, xanthan gum, YE, beef extract, tryptone, casein, keratin, arabinose, xylose, gelatin	0.3–6.0/ 2.5	S ⁰ , Fe (III)	AA, H ₂ , CO ₂	[91]
	<i>Thermosipho atlanticus</i>	Hydrothermal vent, Atlantic Ocean	45–80/ 65	5.0–9.0/ 6.0	1.0–2.6 by 0.2–0.6	cellobiose, xylose, starch, LA, maltose, mannose, trehalose, lactose, arabinose, galactose, mannitol, peptone, casamino acids, gelatin, BHI, YE, glucose	1.5–4.6/ 2.3	S ⁰ , Thio, Cys	AA, iVal, H ₂ , Gly, ALA, Pro	[92]
<i>Geotoga</i>	<i>Geotoga subterranea</i>	Oilfields, USA	30–60/ 45	5.5–9.0/ 6.5	4.0–7.5 by 0.5	mannose, starch, maltodextrins, glucose, lactose, sucrose, galactose, maltose	0.5–10/ 4.0	S ⁰	H ₂ , CO ₂ , AA, EtOH	[10]
	<i>Geotoga petraea</i>	Oilfields, USA	30–55/ 50	5.5–9.0/ 6.5	3.0–20.0 by 0.6	mannose, starch, maltodextrins, glucose, lactose, sucrose, galactose, maltose	0.5–10/ 3.0	S ⁰	H ₂ , CO ₂ , AA, EtOH	[10]
<i>Petrotoga</i>	<i>Petrotoga miotherma</i>	Oilfields, USA	35–65/ 55	5.5–9.0/ 6.5	2.0–7.5 by 0.6	mannose, starch, maltodextrins, glucose, lactose, sucrose, galactose, maltose, maltodextrins, xylose	0.5–10/ 2.0	S ⁰	H ₂ , CO ₂ , AA, EtOH	[10]
	<i>Petrotoga olearia</i>	Oil reservoir, Russia	37–60/ 55	6.5–8.5/ 7.5	0.9–2.5 by 0.3–0.6	arabinose, xylose, cellobiose, dextrin, sucrose, glucose, fructose, maltose, ribose, trehalose, xylan, pyruvate, peptone, starch	0.5–8.0/ 2.0	S ⁰	H ₂ , AA, LA, ALA, EtOH	[93]
	<i>Petrotoga sibirica</i>	Oil reservoir, Russia	37–55/ 55	6.5–9.4/ 8.0	0.9–2.5 by 0.3–0.6	sucrose, glucose, fructose, maltose, ribose, trehalose, xylan, pyruvate, peptone, galactose	0.5–7.0/ 1.0	S ⁰	H ₂ , AA, LA, ALA, EtOH	[93]
<i>Petrotoga</i>	<i>Petrotoga mobilis</i>	Oilfield, North Sea	40–65/ 58–60	5.5–8.5/ 6.5–7.0	1.0–50.0 by 0.5–1.5	starch, xylan, maltodextrin, maltose, cellobiose, sucrose, lactose, glucose, galactose, fructose, arabinose, xylose, ribose, rhamnose	0.5–9.0/ 3.0–4.0	S ⁰ , Thio	H ₂ , CO ₂ , AA, EtOH	[94]
	<i>Petrotoga halophila</i>	Offshore oil, Africa	45–65/ 60	5.6–7.8/ 6.7–7.2	2.0–45.0 by 0.5–0.7	arabinose, cellobiose, fructose, galactose, glucose, lactose, maltose, rhamnose, ribose, starch, sucrose, xylose, xylan, pyruvate	0.5–9.0/ 4.0–6.0	S ⁰	AA, LA, ALA, H ₂ , CO ₂	[95]

Table 1. Cont.

Genus	Species	Isolation	Temp. Range/ Optimal (°C)	pH Range/ Optimal	Cell Dimension (Long by Wide) (µm)	Growth Substrates	NaCl Range/ Optimal (%)	Electron Acceptor	End Products	Ref.
	<i>Petrotoga mexicana</i>	Offshore oil, Africa	25–65/ 55	5.8–8.5/ 6.6	1.0–30.0 by 0.5–0.7	arabinose, cellobiose, fructose, galactose, glucose, lactose, maltose, mannose, raffinose, rhamnose, ribose, starch, sucrose, xylose, xylan, pyruvate.	1.0–20.0/ 3.0	S ⁰ , Thio, Sulfite	AA, LA, H ₂ , CO ₂ , ALA	[96]
	<i>Petrotoga japonica</i>	Oil reservoir, Japan	40–65/ 60	6.0–9.0/ 7.5	2.5–7.0 by 0.25–0.75	starch, xylan, maltose, cellobiose, sucrose, lactose, glucose, galactose, fructose, casamino acids, mannose, arabinose, xylose, ribose	0.5–9.0/ 0.5–1.0	S ⁰ , Thio	AA, H ₂ , CO ₂ , ALA	[97]
	<i>Marinitoga piezophila</i>	Hydrothermal chimney, Pacific Ocean	45–70/ 65	5.0–8.0/ 6.0	1.0–1.5 by 0.5	starch, fructose, glucose, galactose, maltose, cellobiose, ribose, acetate	1.0–5.0/ 3.0	S ⁰ , Thio, Cys	n.d.	[98]
	<i>Marinitoga litoralis</i>	Hot spring, Indian Ocean	45–70/ 65	5.5–7.5/ 6.0	1.0–7.0 by 0.8–1.0	cellobiose, galactose, glucose, glycogen, lactose, maltose, ribose, starch, BHI, casamino acids, casein, peptone, pyruvate, tryptone, YE	0.8–4.6/ 2.6	S ⁰	n.d.	[99]
	<i>Marinitoga okinawensis</i>	Hydrothermal field, Okinawa	30–70/ 55–60	5.5–7.4/ 5.5–5.8	1.5–5.0 by 0.5–0.8	YE, tryptone, peptone, starch, glucose, glycerol	1.0–5.5/ 3.0–3.5	S ⁰ , Cys	n.d.	[100]
	<i>Marinitoga hydrogenitolerans</i>	Hydrothermal chimney, Atlantic Ocean	35–65/ 60	4.5–8.5/ 6.0	1.5–5.0 by 0.5–0.8	glucose, starch, glycogen, chitin, YE, BHI, peptone, casein, pyruvate, maltose	1.0–6.5/ 3.0–4.0	S ⁰ , Thio, Cys	AA, EtOH, Fo, H ₂ , CO ₂	[101]
	<i>Marinitoga artica</i>	Hydrothermal chimney, Norwegian	45–70/ 65	5.0–7.5/ 5.5	1.0–5.0 by 0.5–0.8	glucose, trehalose, maltose, sucrose, maltodextrin, starch, pectin, meat extract, tryptone, YE, pyruvate, fructose, mannose, cellobiose, cellulose, peptone	1.5–5.5/ 2.5	S ⁰ , Cys	n.d.	[102]
	<i>Marinitoga camini</i>	Hydrothermal chimney, Atlantic Ridge	25–65/ 55	5.0–9.0/ 7.0	2.0–3.0 by 0.5–1.0	BHI, gluten, peptone, tryptone, pyruvate, glucose, fructose, maltose, cellobiose, sucrose, starch, cellulose, CMC, pectin, chitin	1.0–4.5/ 2.0	S ⁰ , Cys	AA, iBut, iVal, H ₂ , 3-IAA, LA CO ₂ , HPA, PA	[11]
<i>Oceanotoga</i>	<i>Oceanotoga teriensis</i>	Offshore oil, India	25–70/ 55–58	5.5–9.0/ 7.5	1.5–1.7 by 0.5–0.7	glucose, fructose, cellobiose, arabinose, raffinose, rhamnose, sucrose, xylose, ribose, starch, EtOH, formate, acetate, BHI, YE, bio-tryptidase	0.0–12/ 4.3	S ⁰ , Thio	AA, H ₂ , CO ₂ , EtOH	[12]
<i>Defluviitoga</i>	<i>Defluviitoga tunisiensis</i>	Mesothermic digester	37–65/ 55	6.7–7.9/ 6.9	3.0–30.0 by 1.0	arabinose, cellobiose, fructose, galactose, glucose, lactose, maltose, mannose, raffinose, ribose, sucrose, xylose, cellulose, xylan	0.2–3.0/ 0.5	S ⁰ , Thio	AA, H ₂ , CO ₂	[9]

Table 1. Cont.

Genus	Species	Isolation	Temp. Range/ Optimal (°C)	pH Range/ Optimal	Cell Dimension (Long by Wide) (µm)	Growth Substrates	NaCl Range/ Optimal (%)	Electron Acceptor	End Products	Ref.
<i>Mesotoga</i>	<i>Mesotoga infera</i>	Deep aquifer, France	30–50/ 45	6.2–7.9/ 7.4	2.0–4.0 by 1.0–2.0	arabinose, cellobiose, fructose, galactose, glucose, lactose, LA, mannose, maltose, raffinose, ribose, sucrose, xylose	0.0–1.5/ 0.2	S ⁰	AA, CO ₂	[26]
	<i>Mesotoga prima</i>	Sediment, USA	20–50/ 37	6.5–8.0/ 7.5	1.0 by 0.2	xylose, fructose, ribose, sucrose, mannose, galactose, maltose, lactose, peptone, tryptone, casamino acids, glucose, arabinose, cellobiose, casein, pyruvate	2.0–6.0/ 4.0	S ⁰ , Thio, Sulfite	AA, But, iBut, iVal, 2–MeBu	[8]
<i>Kosmotoga</i>	<i>Kosmotoga arenicorallina</i>	Hot spring, Japan	50–65/ 60	6.2–8.0/ 7.1	1.1–2.7 by 1.1–1.9	xylose, maltose, glycerol	1.0–6.0/ 3.0	S ⁰ , Cys	n.d.	[103]
	<i>Kosmotoga pacifica</i>	Hydrothermal field, Pacific Ocean	33–78/ 70	6.2–8.0/ 7.1	1.0 by 0.6	maltose, YE, peptone, BHI, glycerol, tryptone, xylose, glucose, fructose, cellobiose, trehalose, LA, propionate, glutamate	0.5–6.0/ n.d.	S ⁰ , Cys	n.d.	[104]
	<i>Kosmotoga olearia</i>	Fluid, North Sea	20–80/ 65	5.5–8.0/ 6.8	0.8–1.2 by 0.4–0.7	maltose, ribose, sucrose, starch, casamino acids, tryptone, pyruvate	1.0–6.0/ 2.5–3.0	Thio	H ₂ , CO ₂ , AA, EtOH, PPA	[7]
	<i>Kosmotoga shengliensis</i>	Oilfield, China	45–75/ 65	6.0–8.0/ 7.0	0.7–0.9	glucose, acetate, mEtOH, galactose, fructose, xylose, sucrose, maltose, sorbitol, lactose, xylan, arabinose, formate, rhamnose, glycerol, pyruvate, starch, LA	0.0–4.0/ 1.5	S ⁰ , Thio, Sulfate	AA, LA, ALA, CO ₂ , H ₂	[15]
<i>Athalassatoga</i>	<i>Athalassatoga saccharophila</i>	Hot spring, Japan	30–60/ 55	4.5–7.5/ 5.5–6.0	0.8–2.0 by 0.7–0.8	arabinose, fructose, glucose, lactose, maltose, mannose, ribose, sucrose, xylose, starch, glycogen, peptone, YE	<1/0.0	Fe (III), Thio, Cys	AA, iBut, iVal	[14]
<i>Mesoaciditoga</i>	<i>Mesoaciditoga lauensis</i>	Hydrothermal vent, Pacific Ocean	45–65/ 57–60	4.1–6.0/ 5.5–5.7	0.8–1.0 by 0.4	YE, peptone, maltose, sucrose, glucose, xylose, ribose, starch, tryptone	0.5–6.0/ 3.0	S ⁰ , Thio, Cys	n.d.	[13]

2. Operating Conditions

2.1. H_2 Partial Pressure (P_{H_2})

Since *Thermotogae* members are hydrogen producers, tolerance to hydrogen produced by the bacteria on its own gaseous production, known as the “hydrogen partial pressure (P_{H_2})” effect, is one of the primary parameters being extensively investigated [51,70,105]. The highest hydrogen tolerance has been observed in the genus *Marinotoga*. *Mn. camini* and *Mn. piezophila* were able to grow with H_2 concentrations up to 40% and 60%, respectively. *Mn. hydrogenitolerans* and *Mn. okinawensis* can grow under 100% H_2 atmosphere with only minor inhibition on growth and fermentation [100,101]. Their remarkable resistance to high H_2 levels is probably related to the typical habitats in which *Marinotoga* species thrive [100]. However, the growth of *Thermotogae* species is often inhibited by H_2 accumulation, and the metabolism of these organisms undergoes a series of rearrangements to suit P_{H_2} levels in the bioreactor headspace. The majority of literature data refers to H_2 percentages in gaseous phase, although some studies have been reporting values of P_{H_2} . Partial pressure around 607 mbar led to decreased levels of biomass production, glucose consumption rate, and H_2 production in both *T. neapolitana* and *T. maritima* [106,107]. Boileau et al. [107] highlighted a shift of *T. maritima* glucose catabolism from acetic acid towards lactic acid when P_{H_2} increased from 7 to 607 mbar (Table 2) [106,107]. In contrast, low P_{H_2} (less than 80 mbar) promoted acetic acid accumulation. Biomass production and glucose consumption rate are unaffected when P_{H_2} is maintained within the range of 7.1–178.5 mbar (Table 2) [105,106]. In fact, P_{H_2} lower than 200 mbar is required for optimal growth in reactors, and P_{H_2} around 2900 mbar completely inhibits growth in *T. maritima* [1,45,49,108,109].

Hydrogen evolution is driven by a bifurcating hydrogenase (H_2 ase) that couples the oxidation of reduced ferredoxin (Fd) and NADH with the reduction of protons to H_2 (Figure 1) [58]. In dark fermentation, pyruvate is converted to acetate and ATP, which thermodynamically drives the H_2 -acetate pathway. Under high H_2 partial pressure, hydrogenase activity is inhibited, NADH consumption stops, pyruvate is diverted away from acetic acid production, and lactic acid synthesis becomes the only mechanism for recycling reduced electron carriers (Figure 1) [28–30,57,64,106,110]. Synthesis of lactic acid by the lactate dehydrogenase (LDH) catalyzes the conversion of pyruvate to lactate with the concomitant conversion of NADH to NAD^+ (Figure 1). The depletion of the pyruvate pool, as occurs with the synthesis of lactic acid, negatively affects hydrogen yield, preventing it from reaching the theoretical maximal value (Figure 1) [24]. This problem can be overcome by enhancing the liquid-to-gas mass transfer and keeping H_2 concentrations low in experimental conditions (See Section 2.2) or by using mixed cultures with microbial species that are able to oxidize H_2 [27,111].




Figure 1. Schematic representation of *Thermotogae* metabolic fermentation. Dark fermentation (black arrows) of glucose leads to the production of H₂ and acetate. An increase in CO₂ concentration in the reactor headspace induces the recycling of Ac-CoA and CO₂ into lactate without impairing the synthesis of biogas (blue arrows). This process is named “Capnophilic lactic fermentation (CLF)” [30,31,56,70]. The main end-products of *Thermotogae* fermentation are H₂, lactate, and acetate. Other fermentation products are reported in red. Fe-Fe H₂ase = [Fe-Fe] hydrogenase; PFOR = Pyruvate ferredoxin oxidoreductase; LDH = Lactate dehydrogenase; Fd = Ferredoxin.

2.2. Shaking Speed, Culture/Headspace Volume Ratio, Gas Sparging, and Inoculum

Growth and metabolism of thermophilic bacteria are reported to be strongly affected by an increase in the hydrogen level, which makes the metabolic reactions thermodynamically unfavorable [112]. Many effective strategies have been developed to overcome the H₂ feedback inhibition, such as gas sparging, vigorous stirring, or simply increasing the gas/liquid volume ratio in the reactor. H₂ saturation is dependent on the partial pressure of hydrogen in the culture medium and its mass transfer from liquid to gas phase. As a matter of fact, the mass transfer of H₂ from liquid to gas can be improved by applying vigorous agitation in bioreactors [69,106]. Increased H₂ production rate, glucose consumption rate, and lactic acid synthesis have been observed in *T. neapolitana* cultures with agitation at 200 rpm, compared to static cultures, although the final H₂ yields were similar [106]. Comparable hydrogen yields were also observed when the agitation speed was 300 and 500 rpm, e.g.,

3.0 ± 0.0 mol H₂/mol glucose at 300 rpm vs. 3.2 ± 0.1 mol H₂/mol glucose at 500 rpm, with a mild improvement in fermentation rate (Table 2) [69]. In xylose fermentation, the highest hydrogen and organic acid yields have been reported at 400 rpm when tested in the range of 300–600 rpm [113].

To improve hydrogen liquid-gas mass transfer, Dreschke et al. [69] designed a new method that recirculated the H₂-rich biogas (GaR) into the *T. neapolitana* subs. *capnolactica* broth with agitation (300, 500 rpm). This combination accelerated the H₂ evolution rate and glucose consumption rate during glucose fermentation, compared to the treatments including agitation but excluding GaR. Nonetheless, levels of the end-products, except for H₂ yield, were not significantly altered by the combined parameters (Table 2) [69].

Since P_{H_2} depends on the culture/headspace volume ratio in the bioreactors, its impacts on the performance of fermentation have also been investigated, mainly in batch reactors. Nguyen et al. [64] have experimented various culture/headspace volume ratio from 8.3% (10 mL/120 mL) up to 50% (60 mL/120 mL) in *T. neapolitana* and *T. maritima* cultures [64]. At 8.3%, the H₂ production is the highest for both species (890 mL H₂/L medium in *T. neapolitana* and 883 mL H₂/L medium in *T. maritima*). H₂ production gradually diminished, and lactic acid production was promoted with increasing culture volumes [30,64,110]. d'Ippolito et al. [30] found 1:3 culture/headspace volume was the most suitable ratio for high hydrogen yields [30]. When these conditions were optimized, *T. neapolitana* resulted in H₂ yields between 3.46–3.85 mol H₂/mol glucose [30,114].

Gas sparging, mainly with N₂, is the most common method to reduce hydrogen partial pressure by removing H₂ and CO₂ produced from sugar fermentation in closed bioreactors [56,108,115,116]. Under nitrogen sparging conditions, the overall yield of H₂ in *T. neapolitana* fermentation was about two-fold of the non-sparged cultures, e.g., 1.82 vs. 3.24 mol H₂/mol glucose or 1.14 vs. 2.20 mol H₂/mol xylose (Table 2). The levels of acetic acid and butyrate also increased [110]. Moreover, the fermentation performance was remarkably improved when N₂-sparging was coupled with pH control in *T. neapolitana* using pure glycerol as the sole carbon source (Table 2) [116]. Keeping pH close to neutral improved the glucose utilization and H₂-acetate production rates. In contrast, lactic acid production was lowered under these conditions (0.255 mmol/L with pH control and sparging vs. 0.36 mmol/L with pH control but no sparging) (Table 2) [116]. The use of a CO₂-enriched atmosphere significantly increased both glucose consumption rate and hydrogen production rate, even though the molar yield was comparable to that of N₂-sparging (Table 2) [31]. Surprisingly, supplementation of CO₂ to *T. neapolitana* cultures induced an unexpected metabolic shift from acetic to lactic fermentation without any significant change in hydrogen production (3.6 mol/mol glucose) (Table 2) [31]. Experiments with labeled precursors revealed that part of the exogenous CO₂ was biologically coupled with acetyl-CoA to give lactic acid when the cultures were sparged with CO₂ gas or enriched in sodium bicarbonate (Figure 1) [117]. This process, named Capnophilic Lactic Fermentation (CLF), has the surprising feature to produce more lactic acid than expected from the classical dark fermentation model where H₂ production is impaired by the onset of by-passing pathways (Figure 1) [31,56,117–119]. In dark fermentation, hydrogen and lactic acid levels competed for a common pool of reducing power. Whereas, in CLF, the H₂ level remained high, probably due to additional sources of reductants to sustain NADH-dependent pathways (Figure 1) [118–120]. Recently, an additional increase in lactic acid production occurred in a *T. neapolitana* mutant that was isolated from a culture adapted to continuous exposure to CO₂ [62]. Sparging with CO₂ was also performed on the culture of other *Thermotogales* species, whose metabolic response was qualitatively and quantitatively diverse (Table 2) [70]. CO₂-enriched conditions promoted glucose consumption rate and lowered biogas production in almost all tested species [70]. *T. caldifontis*, *Pseudot. elfii*, *Pseudot. thermarum*, *Pseudot. lettingae*, and *Pseudot. subterranea* did not show substantial variations in the levels of the fermentation products compared to cultures in an N₂-enriched atmosphere [70]. *T. neapolitana*, *T. maritima*, *T. profunda*, and *Pseudot. hypogaea* species responded to CO₂ by reducing the fermentation rate. *T. neapolitana* subs. *capno-*

lactica was the only species to increase lactic acid and H₂ yield moving from N₂-sparging to CO₂-sparging [70]. Generally speaking, the supplementation of external gas (N₂ or CO₂) successfully improves the fermentation performance in most species and lowers the inhibitory effect of H₂ accumulation, but it inevitably causes an undesired dilution of hydrogen in evolved gases. In this context, the recirculation of the H₂-rich biogas method prevents hydrogen saturation in the bioreactor without negatively affecting the content of the produced biogas [69].

The initial biomass concentration (size of inoculum) also has an unexpected impact on the fermentation of thermophilic bacteria. Using various initial biomass concentrations of *T. neapolitana* subs. *capnolactica* (in the range of 0.46–1.74 g CDW/L) under CO₂ atmosphere, hydrogen yield and the distribution of end-products were unaffected (Table 2) [68]. However, increasing inoculum size from 0.46 to 1.74 g/L reduced the fermentation time from 7 h to 3 h [68]. Moreover, the hydrogen production rate, glucose consumption rate, and biomass growth rate were increased [49,50,68]. It is worth pointing out that Ngo et al. [116] reported a reverse correlation between hydrogen production rate and inoculum size, stating that high initial biomass corresponded to a mild reduction of hydrogen production rate [116].

2.3. pH

As the fermentation of sugars leads to the production and accumulation of organic acids, the pH is decreasing during the process, which may inhibit bacterial growth before the substrates are completely consumed [30,106,113]. Two factors impose a strong inhibition on bacterial growth and H₂ production: rapid decrease in pH due to the accumulation of byproducts and feedback inhibition caused by H₂ accumulated in the headspace [65,105–108,113,121].

Thus, pH is a critical factor to control sugar consumption and direct end-products formation [65,67,117,119,122]. Gradual pH drop causes enzyme activity loss [123]. To overcome pH-induced limitations on *Thermotogae* fermentation, several studies were performed with pH adjustments [51,67,121]. In pH-controlled cultures (~6.5–7.0), H₂ and acetic acid production predominated over lactic acid and peaked around 20 h [113]. In contrast, lactic acid production only started when pH declined to around 5.0 [113].

The addition of NaOH at regular intervals and the use of buffering reagents have been regarded as the best-performing methods with serum bottles [56,66,67,113]. The optimum pH for growth and hydrogen production is 6.5–7.0 in *T. maritima* and 6.5–7.5 in *T. neapolitana* depending on substrates and growth conditions [64,113,122]. Moreover, pH 7.0 provides the most promising results in terms of H₂ and organic acids production in *T. neapolitana* [113,122]. A pH shift from 5.5 to 7.0 improved H₂ yield from 125 to 198 mL H₂/L medium in *T. neapolitana* [61]. With *T. neapolitana* cells immobilized on ceramic surfaces using glucose as the carbon source, the highest hydrogen production was observed in the pH range of 7.7–8.5 [51]. Further increase in the range of pH to 8.0–9.0 led to a dramatic decrease in the biogas evolution [64].

Different organic and inorganic buffers have been examined for their effect on anaerobic fermentation under various growth conditions and buffer concentrations [51]. According to Cappelletti et al. [51], 0.1 M HEPES resulted in the best performance, compared to MOPS, PIPES, HPO₄[−]/H₂PO₄[−], or Tris-HCl buffer in *T. neapolitana* batch cultures growing on glucose under N₂ atmosphere [51]. The good buffering properties of HEPES, whose pK (7.55) is near the optimal pH of *T. neapolitana*, was also demonstrated for *T. neapolitana* cultures growing on different complex carbon sources (cheese whey, molasses, or waste glycerol) [51,122]. In another study, 0.05 M HEPES was found to be sufficient under N₂ sparging atmosphere (Table 2) [113]. Under CLF conditions, 0.01 M MOPS, TRIS, or HEPES buffers provided satisfactory results for both H₂ and lactic acid synthesis in *T. neapolitana* subs. *capnolactica* (Table 2) [67]. More specifically, H₂ synthesis was found to be the highest in MOPS, while TRIS promoted acetic acid formation (Table 2) [67]. The highest value of

lactic acid synthesis was 14.9 ± 0.3 mM in phosphate buffer compared to 11.3 ± 0.6 mM in the standard condition (Table 2) [67].

The buffering capacity of HCO_3^- is sufficient to maintain near to optimal pH for growth (~6.5), facilitating the complete substrate degradation and desired by-product formation (Table 2) [31,56,67].

In other studies, itaconic acid was successfully used as a physiological buffer to enhance hydrogen production in *T. neapolitana* growing on glucose or glycerol [121,122]. During the cultivation with 1.5 g/L itaconic acid, the pH slowly dropped from 7.5 to 6.8 over 99 h, while the same pH change was reached within 48 h in cultures not buffered [122]. Although itaconic acid is only poorly catabolized, it affected the overall metabolism of *T. neapolitana* because H_2 and acetic acid production were almost 1.4-fold higher than the control, while lactic acid production was reduced by nearly 100% compared to the control (Table 2) [122]. In addition, Ngo and Sim [122] found that the performance of *T. neapolitana* fermentation growing on waste glycerol was improved by almost 40% by adding itaconic acid into the culture medium [122].

2.4. Temperature

Due to their origin from hot habitats, bacterial species of the phylum *Thermotogae* can live and grow at temperatures in the range of 40–90 °C (Table 1). Some species such as *K. olearia*, *O. teriensis*, *Ms. prima*, and *P. mexicana* can thrive at mesophilic temperatures (Table 1) [7,8,96,100], and other species such as *F. changbaicum*, *F. thailandese*, *T. maritima*, *Pseudot. hypogea*, and *T. neapolitana* share the ability of growing at temperatures close to 90 °C (Table 1) [3,74,77,83,94]. For a long time, researchers have selected an operating temperature of 70 °C [104,117] or 80 °C [105] to cultivate *T. neapolitana* and *T. maritima* without careful investigation of the impacts on fermentation. Nguyen et al. [64] explored changes of H_2 production with temperatures ranging from 55 to 90 °C for *T. neapolitana* and *T. maritima*. Both cultures showed approximately 100 mL H_2 /L medium at 55 °C and a maximum of 200 mL H_2 /L medium at 75–80 °C, with a decrease to 150 H_2 /L medium at 90 °C [64]. In *T. neapolitana*, high temperatures (77–85 °C) enhanced glucose uptake (2.2 mmol/L at 60 °C and 11.0 mmol/L at 77–85 °C) and boosted hydrogen yields (2.04 mol H_2 /mol consumed glucose at 60 °C and 3.85 mol H_2 /mol at 77 °C) [65]. This positive effect was also found for acetic acid (2.0 mmol/L at 60 °C and 18.0 mmol/L at 85 °C) and lactic acid production (no production at 60 °C and 1.25 mmol/L at 85 °C) (Table 2) [65]. Studies conducted on *T. maritima* hydrogenase demonstrated that this enzyme is unstable at the ambient temperature and its activity increased considerably with rising temperature (an activity of 25 units/mg at 20 °C and 110 units/mg at 90 °C [123]).

2.5. Oxygen (O_2)

Thermotogae members occur in various hot ecosystems, including hot springs, deep-sea, and shallow hydrothermal vents, and may also be exposed to O_2 in these ecological niches [1254]. Indeed, despite their anaerobic nature, O_2 tolerance is variable in the phylum; for example, *Thermotoga*, *Fervidobacterium*, and *Geotoga* genera can grow only under strictly anaerobic conditions, while *K. olearia* can survive in up to 15% O_2 [10]. With elemental sulfur, *Ts. atlanticus* can grow with up to 8% O_2 in the headspace [92]. Geochemical and microbial analyses demonstrated the wide distribution of *Thermotogae* species in ecosystems that are not only anaerobic but also partially oxygenated [124]. For this reason, the question of O_2 tolerance and microaerophilic metabolism of *Thermotogae* has been addressed by several studies [65,105,106,125–129]. Some researchers have demonstrated that low concentrations of O_2 are tolerated by *T. neapolitana* and *T. maritima* [127,128]. An O_2 insensitive hydrogenase has been described in *T. neapolitana*, explaining why microaerobic H_2 production and O_2 tolerance could take place in this bacterium [130]. Additionally, *Pseudot. hypogea* and *T. maritima* contain an NADH oxidase that may serve as an O_2 detoxification system [131,132]. Lakhal et al. [129] demonstrated O_2 consumption over 12 h during the stationary phase of *T. maritima* in a batch reactor without reducing agent [129]. O_2 presence

reduced glucose fermentation rate and significantly shifted metabolism towards lactic acid production in *T. maritima* (Table 2). This change can probably be explained by O₂ sensitivity of the hydrogenase [129]. Furthermore, *T. maritima* overproduced enzymes involved in reactive oxygen species (ROS) detoxification, iron-sulfur cluster synthesis/repair, cysteine biosynthesis, and a flavoprotein homologous to the rubredoxin of *Desulfovibrio* species that exhibited an oxygen reductase activity [127].

Van Ooteghem et al. [121] reported that O₂ concentration decreased during the growth of *F. pennavorans*, *P. miotherma*, *Ts. africanus*, *Pseudot. elfii*, and *T. neapolitana*. In these experiments, the H₂ yield greatly exceeded the theoretical limit of 4 mol H₂/mol glucose in *F. pennavorans*, *Pseudot. elfii*, and *T. neapolitana* fermentation [121]. These surprisingly high H₂ yield have led to the hypothesis of an unidentified aerobic pathway using O₂ as a terminal electron acceptor in these bacteria which may not be obligate anaerobes [121]. However, aerobic metabolism is not supported by the genomic sequence of *T. maritima*, although the enzymes involved in the pentose phosphate pathway and an NADPH-reducing hydrogenase have been identified in the genome [16]. To explain the increased yield of H₂ by *T. neapolitana* in microaerobic conditions and the existence of a catabolic process requiring O₂, van Ooteghem et al. [121] used malonic acid as an inhibitor of succinate dehydrogenase and thus the O₂-dependent metabolism. Even if the coding sequence for succinate dehydrogenase has not been identified in the *T. maritima* genome, hydrogen generation was completely inhibited for >40 h in the presence of malonate, postulating that malonate in the medium was no longer available to block catabolism [121]. Then, Eriksen et al. [106] demonstrated that malonic acid was not metabolized by *T. neapolitana* cultures but the exposure to malonic acid clearly affected the metabolism as reduced production of lactic acid and increased H₂ yield were observed [106]. Against these findings, other researchers reported a reduction of H₂ rate and production in *T. neapolitana* cultures after the injection of 6% O₂ [65,106]. The reduction of O₂ consumes reducing equivalents that are then unavailable to produce H₂. The total duration of *T. maritima* fermentation in the batch reactor was delayed about 67 h under O₂-induced stress [129]. In addition, the consumption rate of glucose was drastically reduced and the metabolism of *T. maritima* shifted towards lactic acid production due to inhibition of the O₂-sensitive hydrogenase [129].

From a technical point of view, several strategies were adopted to remove dissolved O₂ in the bioreactor: [I] sparging the culture with N₂, CO₂ or a mixture of both gases; [II] heating the medium; [III] adding a reducing agent such as sodium sulfide or cysteine-HCl in the medium; [IV] maintaining a positive pressure in the bioreactor headspace [31,56,62,67,70,105,106,113,121].

Table 2. Effects of operating conditions on *Thermotoga* fermentation. MOPS: Morpholinopropane-1-sulfonic acid; **HEPES**: 2-[4-(2-hydroxyethyl) piperazin-1-yl] ethanesulfonic acid; **TRIS**: tris(idrossimetil)amminometano cloridrato; **CDW**: Cellular dry weight; **AA**: Acetic acid; **LA**: Lactic acid; **ALA**: Alanine; **But**: Butyrate; **IA**: Itaconic acid; GaR: recirculation of H₂-rich biogas. Experiments were performed in different bioreactor configurations: **B** = Batch; **CSTR** = Continuous-flow Stirred-Tank Reactor; **CSABR**: Continuously Stirred Anaerobic Bioreactor; **SB** = Serum bottles. **H₂ column**: ^a H₂ yield = mol H₂/mol consumed substrate; ^b mL/L culture. * Values extrapolated from the graphical representation of data.

Parameter	Organism	T (°C)	Culture Type	Mixing Speed (rpm)	Reactor/Working Volume (L)	Substrate Loaded (mmol/L)	Operational Parameter	Substrate Consumed (mmol/L)	Products					Ref.
									H ₂ yield ^a	AA (mmol/L)	LA (mmol/L)	ALA (mmol/L)	But (mmol/L)	
<i>P_{H2}</i> (mbar)	<i>T. maritima</i>	80	B	350	1.4/0.1	Glucose (28)	<i>P_{H2}</i> = 7.1 ± 0.4	19.8 ± 1.1	2.34	25.0 ± 1.4	10.5 ± 0.5			[107]
							<i>P_{H2}</i> = 71.4 ± 2.1	19.7 ± 1.4	2.44	24.6 ± 2.4	11.0 ± 0.6			
							<i>P_{H2}</i> = 178.5 ± 3.5	17.2 ± 0.9	2.32	20.1 ± 1.0	9.4 ± 0.5			
							<i>P_{H2}</i> = 606.9 ± 18.7	13.4 ± 0.7	n. d.	13.0 ± 0.7	11.0 ± 0.6			
Stirring Speed (rpm)	<i>T. neapolitana</i>	75	CSABR	300	3.0/1.0	Xylose (33.3)	300	31.43	2.13 ± 0.11	41.8 ± 2.16	1.78 ± 0.11			[113]
				400			400	32.56	2.94 ± 0.15	50.12 ± 2.5	4.0 ± 0.22			
				500			500	32.03	2.31 ± 0.12	44.62 ± 2.16	4.84 ± 0.22			
				600			600	31.87	2.24 ± 0.11	41.12 ± 2.0	1.89 ± 0.11			
	<i>T. neapolitana</i> subsp. <i>capnolactica</i>	80	CSTR	300	3.0/2.0	Glucose (28)	300	22.9 ± 2.7	3.0 ± 0.0	32.3 ± 4.3	10.0 ± 1.0	1.1 ± 0.1		[69]
				500			500	24.8 ± 0.4	3.2 ± 0.1	37.7 ± 2.7	8.1 ± 0.2	1.0 ± 0.1		
				300			300 + GaR	24.7 ± 0.2	3.5 ± 0.2	39.2 ± 1.2	4.4 ± 0.1	0.9 ± 0.0		
				500			500 + GaR	24.9 ± 0.2	3.3 ± 0.1	38.7 ± 2.2	5.1 ± 0.5	0.8 ± 0.0		
Gas sparging	<i>T. neapolitana</i>	80	B	250	3.8/1.0	Glucose (28)	N ₂	25.9 ± 1.3	2.8	44.8 ± 5.4	12.5 ± 2.9	1.3 ± 0.4		[31]
							CO ₂	26.1 ± 1.2	2.8	35.6 ± 5.8	20.0 ± 6.1	2.7 ± 0.5		
		75	SB	no	0.12/0.04	Glycerol (108.6)	w/o	13 ± 0.6	1.24 ± 0.06	8.71 ± 0.35	0.36 ± 0.02			[115]
							N ₂	14 ± 0.7	2.06 ± 0.09	10.04 ± 0.5	0.34 ± 0.02			
Gas sparging	<i>T. neapolitana</i>	77	SB	150	0.12/0.04	Glucose (39)	w/o	-	1.82 ± 0.09	64.28 ± 2.83			33.48 ± 1.47	[110]
							N ₂	-	3.24 ± 0.14	81.42 ± 3.49			36.77 ± 2.04	
						Xylose (27)	w/o	-	1.14 ± 0.07	40.30 ± 3.5			37.68 ± 1.7	
							N ₂	-	2.20 ± 0.13	71.94 ± 3.66			50.62 ± 2.38	
	<i>T. neapolitana</i> subsp. <i>capnolactica</i>	80	SB	no	0.12/0.03	Glucose (28)	N ₂	25.7 ± 0.1	2.5 ± 0.06	27.3 ± 0.8	8.6 ± 0.2	2.5 ± 0.2		[70]
							CO ₂	28.3 ± 1.0	2.9 ± 0.1	22.1 ± 0.9	11.3 ± 0.1	3.0 ± 0.3		

Table 2. Cont.

Parameter	Organism	T (°C)	Culture Type	Mixing Speed (rpm)	Reactor/Working Volume (L)	Substrate Loaded (mmol/L)	Operational Parameter	Substrate Consumed (mmol/L)	Products					Ref.
									H ₂ yield ^a	AA (mmol/L)	LA (mmol/L)	ALA (mmol/L)	But (mmol/L)	
	<i>T. neapolitana</i>	80	SB	no	0.12/0.03	Glucose (28)	N ₂	21.7 ± 0.6	2.5 ± 0.03	30.2 ± 0.4	2.2 ± 0.02	1.9 ± 0.3		[70]
							CO ₂	20.8 ± 2.3	1.9 ± 0.1	20.8 ± 0.1	1.2 ± 0.06	2.4 ± 0.3		
	<i>T. maritima</i>	80	SB	no	0.12/0.03	Glucose (28)	N ₂	23.2 ± 1.0	1.9 ± 0.06	25.5 ± 0.5	5.3 ± 0.8	2.4 ± 0.06		
							CO ₂	19.9 ± 0.6	2.0 ± 0.1	18.3 ± 0.3	1.6 ± 0.2	2.3 ± 0.3		
	<i>T. naphthophila</i>	80	SB	no	0.12/0.04	Glucose (28)	N ₂	13.30 ± 1.10	2.20 ± 0.20	15.70 ± 0.10	1.40 ± 0.06	0.80 ± 0.10		
							CO ₂	20.80 ± 1.70	1.60 ± 0.20	19.20 ± 0.10	5.00 ± 0.02	1.80 ± 0.05		
	<i>T. petrophila</i>	80	SB	no	0.12/0.05	Glucose (28)	N ₂	9.20 ± 1.30	3.00 ± 0.40	13.10 ± 0.05	2.00 ± 0.01	0.00		
							CO ₂	14.20 ± 0.60	1.90 ± 0.10	12.60 ± 0.10	3.80 ± 0.02	0.30 ± 0.10		
	<i>T. caldifontis</i>	70	SB	no	0.12/0.05	Glucose (28)	N ₂	10.90 ± 1.10	2.60 ± 0.10	16.70 ± 3.60	2.20 ± 0.50	3.20 ± 0.90		
							CO ₂	15.20 ± 0.90	1.80 ± 0.03	15.60 ± 1.50	2.30 ± 0.40	6.60 ± 0.70		
	<i>T. profunda</i>	60	SB	no	0.12/0.05	Glucose (28)	N ₂	18.10 ± 0.40	1.50 ± 0.20	15.90 ± 0.40	5.70 ± 0.10	1.40 ± 0.06		
							CO ₂	22.60 ± 1.70	0.70 ± 0.04	5.60 ± 0.20	2.3 ± 0.04	2.60 ± 0.30		
	<i>Pseudot. hypogea</i>	70	SB	no	0.12/0.05	Glucose (28)	N ₂	8.80 ± 1.10	1.10 ± 0.30	6.40 ± 0.10	0.10 ± 0.00	2.90 ± 0.10		
							CO ₂	4.30 ± 0.10	0.50 ± 0.10	3.10 ± 0.20	0.10 ± 0.00	3.40 ± 0.30		
	<i>Pseudot. elfii</i>	70	SB	no	0.12/0.05	Glucose (28)	N ₂	7.00 ± 0.90	2.00 ± 0.20	8.30 ± 0.06	0.20 ± 0.03	4.20 ± 0.30		
							CO ₂	6.70 ± 0.20	2.10 ± 0.10	7.80 ± 0.30	0.10 ± 0.01	10.0 ± 0.30		
	<i>Pseudot. lettingae</i>	70	SB	no	0.12/0.05	Glucose (28)	N ₂	9.30 ± 0.50	1.20 ± 0.10	5.10 ± 0.05	0.20 ± 0.00	2.70 ± 0.05		
							CO ₂	8.10 ± 0.70	1.30 ± 0.30	4.40 ± 0.10	0.05 ± 0.01	3.70 ± 0.20		
Gas sparging	<i>Pseudot. subterranea</i>	70	SB	no	0.12/0.05	Glucose (28)	N ₂	23.10 ± 2.10	1.80 ± 0.20	30.60 ± 6.90	16.20 ± 4.60	9.50 ± 0.40		[70]
							CO ₂	27.00 ± 1.40	1.40 ± 0.10	31.90 ± 7.90	10.70 ± 4.0	20.0 ± 8.0		
	<i>Pseudot. thermarum</i>	80	SB	no	0.12/0.05	Glucose (28)	N ₂	Complete	1.8 ± 0.02	30.00 ± 2.20	6.50 ± 0.20	1.10 ± 0.07		
							CO ₂	Complete	1.50 ± 0.10	24.80 ± 0.70	5.60 ± 0.60	2.20 ± 0.20		
Biomass (g CDW/L)	<i>T. neapolitana</i>	80	Flask	300	0.25/0.2	Glucose (28)	0.46	3.2 ± 0.04	2.39	34.3 ± 0.6	10.9 ± 0.4			[68]
							0.91	2.9 ± 0.06	2.44	32.9 ± 0.8	12.2 ± 0.8			
							1.33	3.4 ± 0.01	2.58	32.3 ± 0.2	11.5 ± 0.5			
							1.74	3.0 ± 0.04	2.37	31.4 ± 1.1	14.7 ± 0.7			

Table 2. Cont.

Parameter	Organism	T (°C)	Culture Type	Mixing Speed (rpm)	Reactor/ Working Volume (L)	Substrate Loaded (mmol/L)	Operational Parameter	Substrate Consumed (mmol/L)	Products					Ref.		
									H ₂ yield ^a	AA (mmol/L)	LA (mmol/L)	ALA (mmol/L)	But (mmol/L)			
pH	<i>T. neapolitana</i> subsp. <i>capnolactica</i>	80	SB	no	0.12/0.03	Glucose (28)	w/o	18.54 ± 0.15	1.78 ± 0.29	22.76 ± 0.40	11.35 ± 0.62	[67]				
							0.01M MOPS	26.42 ± 0.05	3.27 ± 0.18	26.65 ± 0.87	14.23 ± 0.22					
							0.01M TRIS	25.55 ± 0.06	3.10 ± 0.10	26.77 ± 0.29	12.08 ± 0.89					
							0.01M HEPES	25.99 ± 0.03	2.85 ± 0.40	25.56 ± 0.49	13.58 ± 0.88					
							0.01M HCO ₃ [−]	25.62 ± 0.10	2.20 ± 0.30	22.82 ± 0.84	14.63 ± 3.23					
							0.01M phosphate	26.17 ± 0.26	2.78 ± 0.40	24.70 ± 0.59	14.92 ± 0.25					
	<i>T. neapolitana</i>	75	CSABR	300	3.0/1.0	Glucose (28)	w/o pH control	21.98 ± 1.11	2.05 ± 0.1	30.81 ± 1.5	3.33 ± 0.22	[113]				
							plus pH control	27.47 ± 1.39	3.2 ± 0.16	38.3 ± 2.0	1.77 ± 0.11					
						Xylose (33.3)	w/o pH control	29.77 ± 1.46	1.84 ± 0.09	34.47 ± 1.66	3.77 ± 0.22					
							plus pH control	31.83 ± 1.6	2.22 ± 0.11	41.8 ± 2.0	1.66 ± 0.11					
pH	<i>T. neapolitana</i>	75	CSABR	300	3.0/1.0	Sucrose (14.6)	w/o pH control	13.78 ± 0.7	3.52 ± 0.18	33.13 ± 1.65	3.11 ± 0.11	[113]				
							plus pH control	14.69 ± 0.06	4.95 ± 0.25	35.47 ± 1.83	2.11 ± 0.11					
						Xylose (33.3)	w/o pH control	29.44	1.85 ± 0.09	34.97 ± 1.66	3.88 ± 0.22					
							pH = 6.5	32.57	2.71 ± 0.14	49.62 ± 2.50	3.44 ± 0.11					
							pH = 7.0	32.9	2.84 ± 0.14	50.29 ± 2.50	4.00 ± 0.22					
							pH = 7.5	31.77	2.23 ± 0.11	41.96 ± 2.16	1.89 ± 0.11					
						75	SB	no	0.04/ 0.12	Glycerol (108.6)	w/o HEPES		16.96 ± 0.8	1.23 ± 0.06	9.14 ± 0.45	[116]
											0.05 M HEPES		28.26 ± 1.4	2.73 ± 0.14	22.35 ± 1.05	
		pH	<i>T. neapolitana</i>	80	B	250	3.8/1.0	Glucose (28)	w/o NaHCO ₃	25.9 ± 1.3	2.8	44.5 ± 5.4	12.5 ± 2.69	[31]		
									NaHCO ₃ 14 mM	25.4 ± 2.1	1.7	30.5 ± 4.9	18.0 ± 0.6			
NaHCO ₃ 20 mM									23.2 ± 1.9	1.0	44.4 ± 8.2	9.2 ± 2.7				
NaHCO ₃ 40 mM									6.2 ± 0.8	2.7	18.0 ± 4.3	0.7 ± 1.5				
75	B			no	0.12/0.04	Glycerol (108.6)	w/o IA	-	438 ± 22 ^b	7.49 ± 0.33	3.55 ± 0.22 *	[122]				
							1.5 g/L IA	-	619 ± 30 ^b	11.49 ± 0.5	1.66 ± 0.0 *					

Table 2. Cont.

Parameter	Organism	T (°C)	Culture Type	Mixing Speed (rpm)	Reactor/ Working Volume (L)	Substrate Loaded (mmol/L)	Operational Parameter	Substrate Consumed (mmol/L)	Products					Ref.
									H ₂ yield ^a	AA (mmol/L)	LA (mmol/L)	ALA (mmol/L)	But (mmol/L)	
Temp. (°C)	<i>T. neapolitana</i>	60	SB	75	0.26/0.05	Glucose (14)	60	2.2 *	2.04 ± 0.05	2.0	n. d			[65]
		65					65	5.0 *	3.09 ± 0.3	7.0	0.05			
		70					70	8.5 *	3.18 ± 0.02	11.5	0.45			
		77					77	11.0 ± 0.5 *	3.85 ± 0.28	16.5	0.85 ± 0.1			
		85					85	11.0 ± 0.5 *	3.75 ± 0.49	18.0 ± 1.0	1.25 ± 0.05			
Oxygen	<i>T. maritima</i>	80	B	150	2.30/1.53	Glucose (20)	w/o O ₂	17.41	38.09 ^b	18.05	4.36	1.60 ± 0.2		[129]
							with O ₂	19.30	31.75 ^b	18.27	5.45	1.30 ± 0.2		

3. Nitrogen Containing-Compounds

Nitrogen sources (N-sources) are essential for bacterial life for the synthesis of cellular components like nucleic acids, proteins, and enzymes [133,134]. Yeast extract (YE), tryptone, and ammonium chloride (NH_4Cl) have been identified as highly efficient and versatile organic N-sources in laboratory practices. It is widely demonstrated that most of the *Thermotogae* members can use yeast extract and tryptone to grow and metabolize carbohydrates [1,10,77,108,135,136].

Numerous efforts were made to replace YE by combining casamino acids and amino acids, but *Pseudot. elfii* failed to grow on these alternative substrates. The biogas yields of cultures grown with other N-sources were about 4–14% of those with YE (Table 3) [108].

Experiments with different concentrations of YE and tryptone were performed to identify their optimal and minimal concentrations in growth media [64,108,122,137,138]. YE and tryptone are sufficient to ensure growth and hydrogen production without additional carbon sources in *Pseudot. elfii* (Table 3) [108]. van Niel et al. [108] used media with various concentrations of YE and tryptone to ferment glucose by *Pseudot. elfii* [108]. They discovered that increasing the contents of both YE and tryptone from 2 g/L to 5 g/L improved H_2 production (14.8 vs. 28.8 mmol/L) but higher contents did not further improve hydrogen and acetic acid production; high levels of both YE and tryptone only increased acetic acid production in medium lacking other C-sources [108].

When there was a low level of YE (2 g/L) but no tryptone, productions of H_2 and acetic acid remained low, suggesting that tryptone served as an energy source like YE (Table 3) [108]. Although the amino acid compositions of the two N-sources are fairly similar, tryptone contains abundant peptides, a preferred form of amino acids by many bacteria [138]. In another study [122], *T. neapolitana* biomass increased along with the increase of YE concentrations in the range of 1.0–4.0 g/L but not with higher YE concentrations (5.0–6.0 g/L) [122]. The H_2 production plateaued at 420 mL/L in *T. neapolitana* growing on glycerol with 1.0–4.0 g/L YE [122]. Experiments in *T. maritima* and *T. neapolitana* revealed that with over 2 g/L YE, there was a clear increase of acetic acid production, and hydrogen counted up to 30–33% of the total gas in the headspace, even though a mild reduction in glucose consumption occurred (Table 3) [64,138].

Nevertheless, low concentrations (2–4 g/L) of YE are still able to support productivity and bacterial growth [64,108,122,138]. d'Ippolito et al. [30] reported that 2 g/L of both tryptone and YE contributed to 10–15% of the total fermentation products in *T. neapolitana* [30]. Balk et al. [75] demonstrated that *Pseudot. lettingae* was able to degrade methanol in around 30 days in the presence of 0.5 g/L YE, whereas the substrate degradation did not occur when YE was omitted [75]. In contrast, the fermentation of *T. neapolitana* with glucose occurred in a medium without YE, even though the total glucose consumption without YE was attained in 30 h rather than 12 h. H_2 and acetate amounts were half in the medium without YE, (Table 3) [135].

The impact of an inorganic N-source on *Thermotogae* fermentation, such as NH_4Cl , has not been extensively studied, but the presence of NH_4Cl has often been associated with either exopolysaccharide (EPS) formation in *T. maritima* or alanine production in *T. neapolitana* [62,129,136,139]. It is not clear how NH_4Cl stimulates EPS production, but it might involve processing the surplus of reducing equivalents. For example, some organisms produce EPS as a mechanism to transport reducing equivalents out of the cell [140].

Han and Xu [61] demonstrated that a surplus of NH_4Cl could partially substitute YE and tryptone in an optimized medium for auxotrophic *Thermotoga* sp. RQ7 strain [61].

4. Sodium Chloride and Phosphate

All members of the phylum *Thermotogae* showed great adaptability to a wide range of salinity levels (Table 1), although the optimal concentrations of NaCl vary among the members. *Geotoga*, *Oceanotoga*, and *Petrotoga* species can survive in environments comprised of 10% NaCl, while *P. mexicana* can live in up to 20% NaCl (Table 1) [10,12,95].

In contrast, species of the genus *Fervidobacterium* can tolerate salt concentrations up to 1% [5,79–81,83]. Among the species of the genus *Mesotoga*, *Ms. infera* exhibited the lowest tolerance of NaCl (Table 1).

NaCl at 20 g/L was reported to be optimal for *T. neapolitana* growing on either glucose or glycerol when hydrogen production is concerned [64,105,106,108,110,116]. Recently, the effect of different NaCl concentrations (0–35 g/L) on the CLF process was explored in *T. neapolitana* subs. *capnolactica* using glucose as the carbon source [67]. H₂ synthesis and biomass growth were reduced by 15% and 25%, respectively, when NaCl was increased to 35 g/L (Table 3). Similarly, acetic acid production decreased from 26.1 ± 4.7 mM with 10 g/L NaCl to 23.2 ± 0.8 mM with 35 g/L NaCl. In contrast, high NaCl levels had a positive impact on lactic acid production, which increased 7.5-fold (2.8 ± 0.3 mM at 0 g/L NaCl vs. 21.6 ± 6.2 mM at 35 g/L NaCl), without affecting the overall H₂ yields (Table 3) [67]. Pradhan and coworkers [67] suggested a possible involvement of NaCl in a sodium ion gradient that potentially fuels ATP synthesis and transport processes [67]. This creates a bioenergetic balance and supplies necessary reducing equivalents to convert acetic acid into lactic acid under CLF conditions (Figure 1) [67,118,119]. Similarly, another study [141] on H₂-producing *Vibrionaceae* showed that increasing NaCl levels from 9 to 75 g/L enhanced lactic acid synthesis [141].

Regarding phosphate species, they have a strong buffering ability to mitigate pH fluctuation caused by the accumulation of volatile fatty acids [142]. Phosphate deficiency induced an increase in lactic acid production and a small decrease in H₂ formation, suggesting a slight shift of the *T. maritima* metabolism towards lactic acid production. Besides its role as a macro-element, phosphate can also interact with calcium, favoring H₂ production [141,143]. Saidi and co-workers [52] showed that *T. maritima* struggled to produce H₂ at the same rate when there was an oversupply of calcium but an undersupply of phosphate in the medium [52]. For unknown reasons, phosphate exceeding 50 mM has been suggested to inhibit *Pseudot. elfii* growth [108].

Table 3. Effect of organic nitrogen source and NaCl on *Thermotoga* fermentation. **AA:** Acetic acid; **LA:** Lactic acid; **ALA:** Alanine; **YE:** Yeast extract; **Tryp:** Tryptone; **CA:** Casamino acids; **V:** Vitamins solution [108]; **aa:** Amino acids (cysteine, alanine, asparagine, proline, glutamine, serine, and tryptophan, added at 0.2 g/L each). Experiments were performed in different bioreactor configurations: **B** = Batch; **SB** = Serum bottles. **H₂ column:** ^a % H₂ = calculated setting hydrogen production yield on medium with yeast extract to 100%; ^b mmol H₂/L medium; ^c mL H₂/L culture; ^d mol H₂/mol glucose. * Values extrapolated from the graphical representation of data.

Parameter	Organism	T (°C)	Culture Type	Mixing Speed (rpm)	Reactor/ Working Volume (L)	Substrate Loaded (mmol/L)	Operational Parameter	Substrate Consumed (mmol/L)	Products				Ref.
									H ₂	AA (mmol/L)	LA (mmol/L)	ALA (mmol/L)	
Nitrogen sources (g/L)	<i>Pseudot. elfii</i>	65	B	100	3.0/1.0	no	w/o YE	-	40 ^a				[108]
							CA + V	-	4 ^a				
							CA + V + aa	-	6 ^a				
		65	B	100	3.0/1.0	Glucose (22.4)	YE (5)	n.d.	100 ^a				
							CA + V	n.d.	14 ^a				
							CA +V + aa	n.d.	14 ^a				
		65	B	100	3.0/1.0	no	YE (2) -Tryp (0)	-	13.9 ^b	3.5			
							YE (2) -Tryp (2)	-	14.8 ^b	3.4			
							YE (5) -Tryp (0)	-	14.0 ^b	0.0			
							YE (5) -Tryp (5)	-	28.8 ^b	4.9			
		65	B	100	3.0/1.0	Glucose (56)	YE (2) -Tryp (0)	10.3	25.8 ^b	10.7			
							YE (2) -Tryp (2)	18.3	78.5 ^b	19.7			
							YE (5) -Tryp (0)	13.1	84.9 ^b	26.3			
							YE (5) -Tryp (5)	17.9	82.5 ^b	21.2			
	<i>T. neapolitana</i>	80	SB	no	0.12/0.05	Glucose (28)	YE (0.5)	26.6 *	260 ^{ac}	15 *			
							YE (1.0)	26 *	320 ^{ac}	22.5 *			
							YE (2.0)	25.5 *	360 ^{ac}	26.6 *			
							YE (4.0)	25 *	430 ^{ac}	30 *			
							YE (6.0)	25 *	430 ^{ac}	33.3 *			
	<i>T. maritima</i>	80	SB	no	0.12/0.05	Glucose (28.00)	YE (0.5)	25.5 *	190 ^{ac}	0.0 *			
							YE (1.0)	25 *	260 ^{ac}	20.8 *			
YE (2.0)							25 *	270 ^{ac}	23 *				
Nitrogen sources (g/L)	<i>T. maritima</i>	80	SB	no	0.12/0.05	Glucose (28.00)	YE (4.0)	25 *	335 ^{ac}	27.5 *			
							YE (6.0)	24 *	390 ^{ac}	28 *			
	<i>T. neapolitana</i>	77	B	75	0.12/0.05	Glucose (28)	no YE	23 *	9 ^{ab}	4.2 *			
							YE (0.5)	Completed *	16 ^{ab}	7.2 *			

Table 3. Cont.

Parameter	Organism	T (°C)	Culture Type	Mixing Speed (rpm)	Reactor/Working Volume (L)	Substrate Loaded (mmol/L)	Operational Parameter	Substrate Consumed (mmol/L)	Products				Ref.
									H ₂	AA (mmol/L)	LA (mmol/L)	ALA (mmol/L)	
NaCl (g/L)	<i>T. neapolitana</i> subsp. <i>capnolactica</i>	80	SB	no	0.12/0.03	Glucose (28)	w/o	25.62 ± 0.07	2.30 ± 0.50 ^d	20.66 ± 0.27	2.80 ± 0.26	1.28 ± 0.9	[67]
							NaCl (5)	26.00 ± 0.14	2.50 ± 1.20 ^d	24.59 ± 0.95	6.23 ± 3.26	1.61 ± 0.58	
							NaCl (10)	26.12 ± 0.16	3.10 ± 0.80 ^d	26.05 ± 4.69	11.61 ± 2.42	2.46 ± 0.24	
							NaCl (20)	25.96 ± 0.11	3.30 ± 0.20 ^d	25.58 ± 1.03	13.44 ± 0.94	2.41 ± 0.09	
							NaCl (30)	25.68 ± 0.25	2.91 ± 0.37 ^d	23.22 ± 0.81	21.63 ± 6.15	2.38 ± 0.10	

5. Sulfur-Containing Compounds

All members of the phylum *Thermotogae* reduced sulfur-containing compounds such as elemental sulfur (S^0), thiosulfate (Thio), and polysulfide to hydrogen sulfide (H_2S), which is produced at the expense of H_2 (Table 1) [1,4,29,76,144,145]. Sufficient supply of sulfur-containing compounds seems to be critically important; due to a large requirement for Fe-S clusters by the hydrogenase (containing 20 atoms of Fe and 18 atoms of S), PFOR, and other enzymes (Figure 1) [123,146]. In the literature, the effect of sulfur sources has been widely explored. The reduction of S-sources is considered an electron-sink reaction to deplete the surplus of electron power [3,98,107,147]. It is well known that the growth of most anaerobic bacteria of the phylum *Thermotogae* is stimulated by S-sources, but not dependent on them [1,29,52,53,75,107,125,126,144]. Generally speaking, the substrate consumption rate is benefited from a sulfur supply in the medium, except for the methanol fermentation in *Pseudot. lettingae*, which is reduced by S-containing compounds (19.7 mmol/L w/o S-source, 18.7 mmol/L with Thio and 10.6 mmol/L with S^0) (Table 4). Members of the *Mesotoga* genus are able to oxidize sugars, although with low efficiency, only when S^0 is used as the terminal electron acceptor [26,27,66,148,149]. This process gives acetic acid, CO_2 , and sulfide (2 mol of acetate and 4 mol of sulfide per mol of glucose), with no or trace amounts of H_2 (Table 4) [27]. After 250 days of *Ms. prima* cultivation, 9.21 ± 0.13 mmol/L of acetate was measured in the presence of S^0 rather than 1.67 ± 0.21 mM obtained in its absence (Table 4) [27]. Fadhlou and collaborators [27] argued that the metabolic differences between *Thermotoga* spp. and *Ms. prima* strains are related to the absence of a bifurcating [FeFe]-hydrogenase and the accumulation of NADH in *Ms. prima*, leading to growth inhibition in the absence of an external electron acceptor [27]. However, *Ms. prima* and *Ms. infera* strains grew more efficiently in a syntrophic association with a hydrogenotrophic microbial partner that serves as a biological electron acceptor compared to growing *Mesotoga* in a pure culture with sulfur as electron acceptor [26,27]. Boileau et al. [107] investigated the different responses of fermentation performance to different S-sources (Table 4) [107]. Among these compounds (Table 4), thiosulfate, cysteine, and Na_2S were the most efficient ones to optimize *T. maritima* glucose fermentation (Table 4) [107]. Biogas production and glucose utilization increased in the order of no S-source < DMSO < S^0 < Thio < Methionine (Met) < Na_2S < Cysteine (Cys) (Table 4) [107]. Moreover, Na_2S and Cys increased acetic acid production 3-fold and H_2 production 2-fold (Table 4). Thiosulfate seemed to promote lactic acid formation (0.8 ± 0.1 mM w/o S-source and 6.3 ± 0.6 mM with Thio) without affecting other products [107]. Surprisingly, lactic acid was dependent on thiosulfate concentration (0.3 mol/mol glucose w/o Thio and 0.6 mol/mol glucose with 0.24 mmol Thio), even though the proportion between lactic and acetic acid yields remained constant (Table 4). DMSO had no significant impact on *T. maritima* fermentation parameters (Table 4) [107].

In the presence of thiosulfate, the growth and glutamate production of *Fervidobacterium* is stimulated; however, S^0 does not seem to help overcoming the H_2 -feedback inhibition (Table 4) [32,80,88,144]. *P. olearia*, *P. sibirica*, and *Ts. africanus* produced small amounts of ethanol (0.17 mM for both *Petrotoga* species and 0.79 mM for *Ts. africanus*) only in the absence of S-sources (Table 4) [93,145]. *Pseudot. lettingae* produced L-alanine, at the expense of acetic acid, only when thiosulfate or S^0 was present in the medium using methanol as the substrate (Table 4) [75]. Meanwhile, the presence of thiosulfate or S^0 resulted in increased production of acetic acid and decreased production of alanine in *Pseudot. hypogea*, *Ts. melaniensis*, *Ts. geolei*, *P. olearia*, and *P. sibirica* cultures, using glucose or xylose as the carbon source (Table 4) [77,87,90,93]. When S^0 is available, no hydrogen could be detected in *Mn. hydrogenitolerans* growing on glucose [101].

Thermotogae members have been widely employed to degrade different organic wastes, and their degradation significantly benefited from the presence of a reducing agent [51–54,113,116,138]. It is noteworthy to mention that high concentrations of thiosulfinate, a volatile organo-sulfur compound found in organic wastes, has an inhibitory effect on *T. maritima* growth [54]. Similarly, Tao et al. [150] demonstrated that thiosulfinate inhibited the H_2 production by mesophilic seed sludge when co-fermenting food wastes [150].

Table 4. : Effect of sulfur compounds on *Thermotogae* fermentation. **AA**: Acetic acid; **LA**: Lactic acid; **ALA**: Alanine; **EtOH**: Ethanol; **iVal**: isovalerate; **H₂S**: Hydrogen sulfide; **Glu**: Glutamate; **DMSO**: Dimethyl Sulfoxide; **S⁰**: Elemental sulfur; **Met**: Methionine; **Thio**: Thiosulfate; **Cys**: Cysteine; **Na₂S**: Sodium sulfide. * Values extrapolated from the graphical representation of data. ** Concentrations of Sulfur compounds are 0.03 mol equivalent of sulfur. ^a H₂ produced millimolar equivalent; ^b mmol; ^c μM.

Organism	Carbon Source (mM)	Sulfur Source (mM)	Substrate Consumed (mmol/L)	Products mmol/L Culture							Ref.	
				H ₂	AA	LA	ALA	EtOH	iVal	H ₂ S		Glu
<i>T. maritima</i>	Glucose (25)	w/o	7.1 ± 0.4	21.3 ± 2.1	10.1 ± 0.8	0.8 ± 0.1	-					[107]
		DMSO **	9.2 ± 0.5	28.7 ± 2.9	13.3 ± 1.1	0.8 ± 0.1	-					
		S ⁰ **	16.6 ± 0.8	46.1 ± 4.6	23.8 ± 1.9	3.4 ± 0.3	-					
		Met **	18.3 ± 0.9	53.3 ± 5.3	26.5 ± 2.1	3.1 ± 0.3	-					
		Thio **	17.5 ± 0.9	47.3 ± 4.7	24.1 ± 1.9	6.3 ± 0.6	-					
		Cys **	20.4 ± 1.0	58.5 ± 5.8	30.5 ± 2.4	4.1 ± 0.4	-					
		Na ₂ S **	20.4 ± 1.0	54.9 ± 5.5	30.7 ± 2.5	4.7 ± 0.5	-					
	Glucose (60)	w/o Thio	17.7 ± 1.9	25.0 ± 2.2	12.8 ± 1.0	5.4 ± 0.6	1.39 ± 0.2					
		Thio (0.01)	20.0 ± 1.1	31.0 ± 2.3	16.0 ± 0.8	10.2 ± 1.1	-					
		Thio (0.03)	28.0 ± 1.5	57.9 ± 4.8	30.6 ± 1.9	8.2 ± 0.7	-					
		Thio (0.06)	38.5 ± 2.0	73.3 ± 5.9	38.2 ± 2.4	18.1 ± 1.8	-					
		Thio (0.12)	45.7 ± 2.5	99.7 ± 8.3	52.4 ± 3.3	15.4 ± 1.6	3.8 ± 0.3					
		Thio (0.18)	45.4 ± 2.2	86.9 ± 8.2	45.0 ± 2.2	23.4 ± 2.3	-					
		Thio (0.24)	43.8 ± 2.2	88.6 ± 8.9	46.1 ± 3.3	26.4 ± 1.4	3.8 ± 0.2					
Glucose (20)	w/o	13.70	36.09	15.62		0.70			n.d.		[145]	
	Thio (20)	13.55	4.02	15.99		0.80			14.45			
<i>T. neapolitana</i>	Glucose (20)	w/o	14.00	31.67	18.27		0.87			n.d.		[145]
		Thio (20)	13.90	16.07	16.12		0.60			7.39		
<i>Pseudot. lettingae</i>	Methanol (20)	w/o	19.70	n. d.	13.70		-			-		[75]
		Thio (20)	18.7	n. d.	-		5.8			11.2		
		S ⁰ (2%)	10.6	n. d.	-		3.1			7.3		
<i>Pseudot. hypogea</i>	Glucose (20)	w/o	8.60	29.03	4.49		1.71			n. d.		[145]
		Thio (20)	14.39	2.29	19.7		1.06			15.08		

Table 4. Cont.

Organism	Carbon Source (mM)	Sulfur Source (mM)	Substrate Consumed (mmol/L)	Products mmol/L Culture								Ref.
				H ₂	AA	LA	ALA	EtOH	iVal	H ₂ S	Glu	
<i>Pseudot. hypogea</i>	Glucose (20)	w/o	7.0	9.4 ^a	5.0		1.7	1.0		0.2		[77]
		Thio (20)	13.0	0.9 ^a	19.8		1.0	1.6		15.1		
<i>Pseudot. hypogea</i>	Xylose (20)	w/o	12.9	19.0 ^a	8.9		2.4	1.0		0.2		[77]
		Thio (20)	12.0	1.8 ^a	13.7		1.3	1.0		7.5		
<i>Pseudot. elfii</i>	Glucose (20)	w/o	3.1	8.8	4.0					0.0		[77]
		Thio (20)	10.4	2.0	17.9					23.00		
	Glucose (20)	w/o	2.75	7.70	3.49		1.05			n. d.		[145]
		Thio (20)	8.15	n. d.	12.63		0.41			14.55		
<i>Ts. geolei</i>	Glucose (0.28)	w/o	7.0 ^b	9.3 ^a	8.5 ^b		1.2 ^b			0.5 ^b		[87]
		S ⁰ (2%)	6.0 ^b	0.0 ^a	7.5 ^b		0.5 ^b			12.5 ^b		
<i>Ms. Prima Phos Ac3</i>	Glucose (20)	w/o	1.50 ± 0.20	<1 ^c	1.67 ± 0.21					1.05 ± 0.25		[27]
		S ⁰	6.57 ± 0.19	<1 ^c	9.21 ± 0.13					24.40 ± 0.30		
<i>Ms. Prima MesGIAg4.2T</i>	Fructose (20)	w/o	1.00 ± 0.23	<1 ^c	0.70 ± 0.41					1.18 ± 0.41		
		S ⁰	3.27 ± 0.85	<1 ^c	8.48 ± 1.96					18.03 ± 5.16		
<i>Ts. africanus</i>	Glucose (28)	w/o	7.20	16.80	7.90	<0.2		0.79		n.d.		[145]
		Thio (20)	7.70	1.00	12.40	-		-		14.60		
<i>Ts. atlanticus</i>	Glucose (28)	w/o	5.6	12.5	1.7				0.14	-		[92]
		S ⁰ (1%)	6.0	7.5	1.9				0.15	1.3		
<i>F. islandicum</i>	Glucose (20)	w/o	14.20	21.58	6.25		3.98			n.d.		[145]
		Thio (20)	16.20	n. d.	20.25		1.22			34.02		
<i>F. pennavorans</i>	Glucose (11)	w/o	-	0.25 *	6.7 *		4.0 ± 0.5 *				1.3 *	[32]
		Thio (20)	-	0.2 *	6.7 *		4.50 *				No *	

6. Metal Ions

Typically, hydrothermal ecosystems are enriched with essential micronutrients and trace metals such as soluble and insoluble iron, manganese, cobalt, and molybdenum. Some terrestrial hydrothermal waters are also characterized by chromium and uranium contents of several micrograms per liter [151]. The physiological roles that most of these metals play in microbial metabolism are still largely unknown. It is believed that their functions include energy generation and biosynthesis [151]. In addition, Mn, Fe, Zn, and Co metals are vitally important micro-elements for growth, essential for cellular transport processes, and serve as cofactors for many enzymes [152]. Understanding the physicochemical properties of extreme habitats can help to determine the metal toxicity limits on microbial growth in laboratory settings. Indeed, metal susceptibility tests have been carried out on *T. neapolitana*, *T. maritima*, and *Ts. africanus*, and have identified the following toxicity order: cadmium (1.0–10.0 μM) > zinc (0.01–0.1 mM) > nickel (1.0–5.0 mM) > cobalt (1.0–10.0 mM) [153].

Attention has also been paid to Fe (III) reduction by thermophilic bacteria, since Fe (III) may work as an external electron acceptor in microbial metabolism [154]. Members of the phylum *Thermotogae* are capable of coupling the reduction of iron with the oxidation of a wide range of organic and inorganic compounds. *T. maritima* reduced Fe (III) into Fe (II) exclusively with molecular hydrogen as an electron donor [154]. Fe (III) reduction has also been reported to stimulate growth and mitigate H_2 inhibition in *Pseudot. lettingae*, *Pseudot. subterranea*, *Pseudot. elfii*, *Ts. affectus*, *Ts. globiformans*, and *Ts. activus* [75,76,88,89,91]. The recently characterized member of the order *Mesoaciditogales*, *A. saccharophila*, changed fermentation end-products when growing with Fe (III), favoring the production of small amounts of acetate, isobutyrate, and isovalerate [14].

Ions and metals are generally supplied in *Thermotogae* growth media through Balch's oligo-elements solution [155]. The removal of oligo-elements from *T. maritima* cultures resulted in a minor increase in lactic acid production (1.2 vs. 4.3 mmol/L) and a decrease in H_2 productivity (12.4 vs. 8.8 mmol/h/L) [52]. Limitation in iron lowered H_2 production by deviating the fermentation pathway towards the production of more reduced end-products such as lactic acid in mixed cultures [156,157]. Another study [139] highlighted how the supplementation of Fe ions to mixed cultures had pronounced effect on hydrogen activity [139]. Similarly, Fe^{2+} (as well as Co, Ni and Mn) stimulated *Pseudot. hypogea* alcohol dehydrogenase activity (ADH), an iron-containing enzyme involved in alcohol fermentation, by 10–15%, while Zn^{2+} completely inhibited the enzyme activity [158]. On the same base, the inclusion of tungsten in the growth medium of *T. maritima* increased the specific activity of both hydrogenase (by up to 10-fold) and PFOR in cell-free extracts, although the function of tungsten in the metabolism of *T. maritima* is not clear [123,126].

As for magnesium, potassium, and calcium ions, they not only play critical roles in bacterial growth, but also act as enzyme cofactors and ensure the survival of microorganisms in their hot ecosystems, by protecting double-stranded DNA from degradation [159]. The best cell yields were obtained with a low concentration of Mg^{2+} and a high concentration of Ca^{2+} [126]. It would be worthwhile to dig further into the metal ions repercussions on *Thermotogae* metabolism in future research.

7. Conclusions

Steam reforming of methane (CH_4) is currently used to produce hydrogen in the industry, as it is the most economic technology available so far. Producing hydrogen by biological means at an industrial scale remains as a challenge. Within the race to find the best way to generate hydrogen via microbes (e.g., choice of strains, substrates, fermentation conditions), *Thermotogae* seem to have many unique advantages. Optimization of their cultivation conditions is fundamental to improve the overall productivity of the fermentation system and its profitability, which determine the feasibility of replacing the current methods of hydrogen production.

The phylum *Thermotogae* comprises a wide collection of species with astonishing and unique features associated to their original habitats. Extensive research has shown tremen-

dous potentials of using these bacteria in biological production of hydrogen, degradation of wastes, and isolation of thermostable enzymes.

Many factors affect the anaerobic metabolism of *Thermotogae* species, including operating conditions (shaking, inoculum, gas sparging, and culture/headspace volume ratio), temperature, pH, nitrogen, sulfur-containing compounds, sodium chloride, phosphate, and metal ions. Optimization of these fermentation parameters has been intensively pursued with *Thermotoga* and *Pseudothermotoga* species, which are the best hydrogen producers in the phylum. In contrast, little is known regarding other species of the phylum, especially their ability to synthesize desirable biological products.

In general, *Thermotogae* fermentation is affected by the accumulation of produced biogas and organic acids because they increase hydrogen partial pressure inside of the bioreactor and drastically reduce the pH of the cultivation medium. Consequently, the metabolic process stops before the substrate is completely consumed. Gas sparging, stirring, and adjusting culture/headspace volume ratio can help to overcome the inhibition on growth caused by hydrogen accumulation. Implementing these strategies and adjusting pH during the fermentation process can result in high hydrogen yields and efficient consumption of substrates. A reduction of fermentation time by starting with the right inoculum size could cast favorable great perspectives on the economics of the industrial processes.

This review highlights the importance of nitrogen-containing compounds that need to be supplied to the medium to stimulate bacterial growth. Overall, yeast extract and tryptone are the preferred forms of nitrogen. Sulfur-containing compounds not only play a critical role in bacterial growth but also divert reducing power to selectively produce certain end-products in *Thermotogae* metabolism.

Until now, the impact of metal ions and salts on the fermentation process has not been well investigated even though it has been demonstrated that they could stimulate many key enzymes involved in various metabolic pathways.

In summary, the extensive data collection of this review offers a great reference for the optimization and development of sustainable bioprocesses based on *Thermotogae* species and helps to generate insightful perspectives for the exploitation of these anaerobic bacteria in biotechnological processes.

Funding: This research was funded by BioRECO₂VER Project, through the European Union's Horizon 2020 Research and Innovation Programme under Grant Agreement No. 760431.

Acknowledgments: The authors would like to thank Lucio Caso (CNR-ICB) for the technical support in preparing the manuscript.

Conflicts of Interest: The authors declare no conflict of interest. The funding agencies gave their permissions to the publication of manuscript.

References

1. Huber, R.H.M. *Thermotogales*; Dworkin, M., Falkow, S., Rosenberg, E., Schleifer, K.H., Stackebrandt, E., Eds.; Springer: New York, NY, USA, 2006; pp. 899–922.
2. Bhandari, V.; Gupta, R.S. The Phylum Thermotogae. In *The Prokaryotes*; Rosenberg, E., DeLong, E.F., Lory, S., Stackebrandt, E., Thompson, F., Eds.; Springer: Berlin/Heidelberg, Germany, 2014.
3. Huber, R.; Langworthy, T.A.; König, H.; Thomm, M.; Woese, C.R.; Sleytr, U.B.; Stetter, K.O. *Thermotoga maritima* Sp. Nov. Represents a New Genus of Unique Extremely Thermophilic Eubacteria Growing up to 90 °C. *Arch. Microbiol.* **1986**, *144*, 324–333. [[CrossRef](#)]
4. Belahbib, H.; Summers, Z.M.; Fardeau, M.; Joseph, M.; Tamburini, C.; Dolla, A.; Ollivier, B.; Armougom, F. Towards a Congruent Reclassification and Nomenclature of the Thermophilic Species of the Genus *Pseudothermotoga* within the Order Thermotogales. *Syst. Appl. Microbiol.* **2018**, *41*, 555–563. [[CrossRef](#)] [[PubMed](#)]
5. Patel, B.K.C.; Morgan, H.W.; Daniel, R.M. *Fervidobacterium nodosum* Gen. Nov. and Spec. Nov., a New Chemoorganotrophic, Caldoactive, Anaerobic Bacterium. *Arch. Microbiol.* **1985**, *141*, 63–69. [[CrossRef](#)]
6. Windberger, E.; Huber, R.; Trincone, A.; Fricke, H.; Stetter, K.O. *Thermotoga thermarum* Sp. Nov. and *Thermotoga Neapolitana* Occurring in African Continental Solfataric Springs. *Arch. Microbiol.* **1989**, *151*, 506–512. [[CrossRef](#)]

7. DiPippo, J.L.; Nesbø, C.L.; Dahle, H.; Doolittle, W.F.; Birkland, N.K.; Noll, K.M. *Kosmotoga olearia* Gen. Nov., Sp. Nov., a Thermophilic, Anaerobic Heterotroph Isolated from an Oil Production Fluid. *Int. J. Syst. Evol. Microbiol.* **2009**, *59*, 2991–3000. [[CrossRef](#)] [[PubMed](#)]
8. Nesbø, C.L.; Bradnan, D.M.; Adebuseyi, A.; Dłutek, M.; Petrus, A.K.; Foght, J.; Doolittle, W.F.; Noll, K.M. *Mesotoga prima* Gen. Nov., Sp. Nov., the First Described Mesophilic Species of the Thermotogales. *Extremophiles* **2012**, *16*, 387–393. [[CrossRef](#)]
9. Ben Hania, W.; Godbane, R.; Postec, A.; Hamdi, M.; Ollivier, B.; Fardeau, M.L. *Defluviitoga tunisiensis* Gen. Nov., Sp. Nov., a Thermophilic Bacterium Isolated from a Mesothermic and Anaerobic Whey Digester. *Int. J. Syst. Evol. Microbiol.* **2012**, *62*, 1377–1382. [[CrossRef](#)]
10. Davey, M.E.; Wood, W.A.; Key, R.; Nakamura, K.; Stahl, D.A. Isolation of Three Species of Geotoga and Petrotoga: Two New Genera, Representing a New Lineage in the Bacterial Line of Descent Distantly Related to the “Thermotogales”. *Syst. Appl. Microbiol.* **1993**, *16*, 191–200. [[CrossRef](#)]
11. Wery, N.; Lesongeur, F.; Pignet, P.; Derennes, V.; Cambon-Bonavita, M.A.; Godfroy, A.; Barbier, G. *Marinitoga camini* Gen. Nov., Sp. Nov., a Rod-Shaped Bacterium Belonging to the Order Thermotogales, Isolated from a Deep-Sea Hydrothermal Vent. *Int. J. Syst. Evol. Microbiol.* **2001**, *51*, 495–504. [[CrossRef](#)]
12. Jayasinghearachchi, H.S.; Lal, B. *Oceanotoga Teriensis* Gen. Nov., Sp. Nov., a Thermophilic Bacterium Isolated from Offshore Oil-Producing Wells. *Int. J. Syst. Evol. Microbiol.* **2011**, *61*, 554–560. [[CrossRef](#)]
13. Reysenbach, A.L.; Liu, Y.; Lindgren, A.R.; Wagner, I.D.; Sislak, C.D.; Mets, A.; Schouten, S. *Mesoaciditoga lauensis* Gen. Nov., Sp. Nov., a Moderately Thermoacidophilic Member of the Order Thermotogales from a Deep-Sea Hydrothermal Vent. *Int. J. Syst. Evol. Microbiol.* **2013**, *63*, 4724–4729. [[CrossRef](#)] [[PubMed](#)]
14. Itoh, T.; Onishi, M.; Kato, S.; Iino, T.; Sakamoto, M.; Kudo, T.; Takashina, T.; Ohkuma, M. *Athalassotoga saccharophila* Gen. Nov., Sp. Nov., Isolated from an Acidic Terrestrial Hot Spring, and Proposal of Mesoaciditogales Ord. Nov. and Mesoaciditogaceae Fam. Nov. in the Phylum Thermotogae. *Int. J. Syst. Evol. Microbiol.* **2016**, *66*, 1045–1051. [[CrossRef](#)] [[PubMed](#)]
15. Feng, Y.; Cheng, L.; Zhang, X.; Li, X.; Deng, Y.; Zhang, H. *Thermococcoides shengliensis* Gen. Nov., Sp. Nov., a New Member of the Order Thermotogales Isolated from Oil-Production Fluid. *Int. J. Syst. Evol. Microbiol.* **2010**, *60*, 932–937. [[CrossRef](#)] [[PubMed](#)]
16. Nelson, K.; Clayton, R.; Gill, S.; Gwinn, M.; Dodson, R.; Haft, D.; Hickey, E.; Peterson, J.; Nelson, W.; Ketchum, K.; et al. Evidence for Lateral Gene Transfer between Archae and Bacteria from Genome Sequence of *Thermotoga maritima*. *Nature* **1999**, *399*, 323–329. [[CrossRef](#)] [[PubMed](#)]
17. Connors, S.B.; Mongodin, E.F.; Johnson, M.R.; Montero, C.I.; Nelson, K.E.; Kelly, R.M. Microbial Biochemistry, Physiology, and Biotechnology of Hyperthermophilic Thermotoga Species. *FEMS Microbiol. Rev.* **2006**, *30*, 872–905. [[CrossRef](#)]
18. Rodionov, D.A.; Rodionova, I.A.; Li, X.; Ravcheev, I.; Tarasova, Y.; Portnoy, V.A.; Zengler, K.; Osterman, A.L. Transcriptional Regulation of the Carbohydrate Utilization Network in *Thermotoga maritima*. *Front. Microbiol.* **2013**, *4*, 244. [[CrossRef](#)]
19. Galperin, M.Y.; Noll, K.M.; Romano, A.H. The Glucose Transport System of the Hyperthermophilic Anaerobic Bacterium *Thermotoga neapolitana*. *Appl. Environ. Microbiol.* **1996**, *62*, 2915–2918. [[CrossRef](#)] [[PubMed](#)]
20. Paulsen, I.T.; Nguyen, L.; Sliwinski, M.K.; Rabus, R.; Jr, M.H.S. Microbial Genome Analyses: Comparative Transport Capabilities in Eighteen Prokaryotes. *J. Mol. Biol.* **2000**, *301*, 75–100. [[CrossRef](#)]
21. Nanavati, D.; Thirangoon, K.; Noll, K.M. Several Archaeal Homologs of Putative Oligopeptide-Binding Proteins Encoded by *Thermotoga maritima* Bind Sugars. *Appl. Environ. Microbiol.* **2006**, *72*, 1336–1345. [[CrossRef](#)]
22. Latif, H.; Sahin, M.; Tarasova, J.; Tarasova, Y.; Portnoy, V.A.; Nogales, J.; Zengler, K. Adaptive Evolution of *Thermotoga maritima* Reveals Plasticity of the ABC Transporter Network. *Appl. Environ. Microbiol.* **2015**, *81*, 5477–5485. [[CrossRef](#)]
23. Boucher, N.; Noll, K.M. Substrate Adaptabilities of Thermotogae Mannan Binding Proteins as a Function of Their Evolutionary Histories. *Extremophiles* **2016**, *20*, 771–783. [[CrossRef](#)] [[PubMed](#)]
24. Thauer, R.K.; Jungermann, K.; Decker, K. Energy Conservation in Chemotrophic Anaerobic Bacteria. *Bacteriol. Rev.* **1977**, *41*, 100–180. [[CrossRef](#)] [[PubMed](#)]
25. Ben Hania, W.; Ghodbane, R.; Postec, A.; Brochier-Armanet, C.; Hamdi, M.; Fardeau, M.L.; Ollivier, B. Cultivation of the First Mesophilic Representative (“Mesotoga”) within the Order Thermotogales. *Syst. Appl. Microbiol.* **2011**, *34*, 581–585. [[CrossRef](#)] [[PubMed](#)]
26. Hania, W.B.; Postec, A.; Aüllo, T.; Ranchou-Peyruse, A.; Erauso, G.; Brochier-Armanet, C.; Hamdi, M.; Ollivier, B.; Saint-Laurent, S.; Magot, M.; et al. *Mesotoga infera* Sp. Nov., a Mesophilic Member of the Order Thermotogales, Isolated from an Underground Gas Storage Aquifer. *Int. J. Syst. Evol. Microbiol.* **2013**, *63*, 3003–3008. [[CrossRef](#)] [[PubMed](#)]
27. Fadhlou, K.; Hania, W.B.; Armougom, F.; Bartoli, M.; Fardeau, M.L.; Erauso, G.; Brasseur, G.; Aubert, C.; Hamdi, M.; Brochier-Armanet, C.; et al. Obligate Sugar Oxidation in Mesotoga Spp., Phylum Thermotogae, in the Presence of Either Elemental Sulfur or Hydrogenotrophic Sulfate-Reducers as Electron Acceptor. *Environ. Microbiol.* **2017**, *20*, 281–292. [[CrossRef](#)] [[PubMed](#)]
28. Janssen, P.H.; Morgan, H.W. Heterotrophic Sulfur Reduction by Thermotoga Sp. Strain FjSS3.B1. *FEMS Microbiol. Lett.* **1992**, *96*, 213–218. [[CrossRef](#)] [[PubMed](#)]
29. Schröder, C.; Selig, M.; Schönheit, P. Glucose Fermentation to Acetate, CO₂ and H₂ in the Anaerobic Hyperthermophilic Eubacterium *Thermotoga maritima*: Involvement of the Embden-Meyerhof Pathway. *Arch. Microbiol.* **1994**, *161*, 460–470. [[CrossRef](#)]
30. D’Ippolito, G.; Dipasquale, L.; Vella, F.M.; Romano, I.; Gambacorta, A.; Cutignano, A.; Fontana, A. Hydrogen Metabolism in the Extreme Thermophile *Thermotoga neapolitana*. *Int. J. Hydrogen Energy* **2010**, *35*, 2290–2295. [[CrossRef](#)]

31. Dipasquale, L.; d'Ippolito, G.; Fontana, A. Capnophilic Lactic Fermentation and Hydrogen Synthesis by *Thermotoga neapolitana*: An Unexpected Deviation from the Dark Fermentation Model. *Int. J. Hydrogen Energy* **2014**, *39*, 4857–4862. [\[CrossRef\]](#)
32. Wushke, S.; Fristensky, B.; Zhang, X.L.; Spicer, V.; Krokhin, O.V.; Levin, D.B.; Stott, M.B.; Sparling, R. A Metabolic and Genomic Assessment of Sugar Fermentation Profiles of the Thermophilic Thermotogales, *Fervidobacterium pennivorans*. *Extremophiles* **2018**, *22*, 965–974. [\[CrossRef\]](#)
33. Vijayakumar, J.; Aravindan, R.; Viruthagiri, T. Lactic Acid and Its Potential Applications in Industries. *Bioprod. Biosyst. Eng.* **2007**, *42*, 101–103.
34. Roehr, M.; Kosaric, N.; Vardar-Sukan, F.; Pieper, H.J.; Senn, T. *The Biotechnology of Ethanol. Classical and Future Applications*; Wiley: Hoboken, NJ, USA, 2005; p. 245. ISBN 978-3-527-60234-6.
35. Vieille, C.; Zeikus, G.J. Hyperthermophilic Enzymes: Sources, Uses, and Molecular Mechanisms for Thermostability. *Microbiol. Mol. Biol. Rev.* **2001**, *65*, 1–43. [\[CrossRef\]](#) [\[PubMed\]](#)
36. Ul Haq, I.; Hussain, Z.; Khan, M.; Muneer, B.; Afzal, S.; Majeed, S.; Akram, F. Kinetic and Thermodynamic Study of Cloned Thermostable Endo-1, 4- β -Xylanase from *Thermotoga petrophila* in Mesophilic Host. *Mol. Biol. Rep.* **2012**, *39*, 7251–7261. [\[CrossRef\]](#) [\[PubMed\]](#)
37. Ul Haq, I.; Tahir, S.F.; Aftab, M.N.; Akram, F.; ur Rehman, A.; Nawaz, A.; Mukhtar, H. Purification and Characterization of a Thermostable Cellobiohydrolase from *Thermotoga petrophila*. *Protein Pept. Lett.* **2018**, *25*, 1003–1014. [\[CrossRef\]](#) [\[PubMed\]](#)
38. Colussi, F.; Viviam, M.; Ian, S.; Junio, M. Oligomeric State and Structural Stability of Two Hyperthermophilic β -Glucosidases from *Thermotoga petrophila*. *Amino Acids* **2015**, *47*, 937–948. [\[CrossRef\]](#) [\[PubMed\]](#)
39. Pollo, S.M.J.; Zhaxybayeva, O.; Nesbø, C.L. Insights into Thermoadaptation and the Evolution of Mesophily from the Bacterial Phylum Thermotogae. *Can. J. Microbiol.* **2015**, *61*, 655–670. [\[CrossRef\]](#) [\[PubMed\]](#)
40. Fatima, B.; Aftab, M.; Ul Haq, I. Cloning, Purification, and Characterization of Xylose Isomerase from *Thermotoga naphthophila* RKU-10. *J. Basic Microbiol.* **2016**, *56*, 949–962. [\[CrossRef\]](#)
41. Lopes, J.L.S.; Yoneda, J.S.; Martins, J.M.; Demarco, R. Environmental Factors Modulating the Stability and Enzymatic Activity of the *Petrotoga mobilis* Esterase (PmEst). *PLoS ONE* **2016**, *11*, e0158146. [\[CrossRef\]](#)
42. Intagun, W.; Kanoksilapatham, W. A Review: Biodegradation and Applications of Keratin Degrading Microorganisms and Keratinolytic Enzymes, Focusing on Thermophiles and Thermostable Serine Proteases. *Am. J. Appl. Sci. Rev.* **2017**, *14*, 1016–1023. [\[CrossRef\]](#)
43. Hamid, A.; Aftab, M.N. Cloning, Purification, and Characterization of Recombinant Thermostable β -Xylanase Tnap_0700 from *Thermotoga naphthophila*. *Appl. Biochem. Biotechnol.* **2019**, *189*, 1274–1290. [\[CrossRef\]](#)
44. Kang, E.; Jin, H.; La, J.W.; Park, S.; Kim, W.; Lee, W. Identification of Keratinases from *Fervidobacterium islandicum* AW-1 Using Dynamic Gene Expression pro Fi Ling. *Microb. Biotechnol.* **2019**, *13*, 442–457. [\[CrossRef\]](#) [\[PubMed\]](#)
45. European Commission. *Communication from the Commission to the European Parliament, the Council, the European Economic and Social Committee and the Committee of the Regions*; A Hydrogen Strategy for a Climate-Neutral Europe; 8.7.2020 COM 301 Final; European Commission: Brussels, Belgium, 2020.
46. Dipasquale, L.; d'Ippolito, G.; Gallo, C.; Vella, F.M.; Gambacorta, A.; Picariello, G.; Fontana, A. Hydrogen Production by the Thermophilic Eubacterium *Thermotoga neapolitana* from Storage Polysaccharides of the CO₂-Fixing Diatom *Thalassiosira weissflogii*. *Int. J. Hydrogen Energy* **2012**, *37*, 12250–12257. [\[CrossRef\]](#)
47. Angenent, L.; Karim, K.; Al-Dahhan, M.; Wrenn, B.; Domiguez-Espinosa, R. Production of Bioenergy and Biochemicals from Industrial and Agricultural Wastewater. *Trends Biotechnol.* **2004**, *22*, 477–485. [\[CrossRef\]](#) [\[PubMed\]](#)
48. De Vrije, T.; de Haas, G.G.; Tan, G.B.; Keijsers, E.R.P.; Claassen, P.A.M. Pretreatment of Miscanthus for Hydrogen Production by *Thermotoga elfii*. *Hydrog. Energy* **2002**, *27*, 1381–1390. [\[CrossRef\]](#)
49. De Vrije, T.; Budde, M.A.W.; Lips, S.J.; Bakker, R.R.; Mars, A.E.; Claassen, P.A.M. Hydrogen Production from Carrot Pulp by the Extreme Thermophiles *Caldicellulosiruptor saccharolyticus* and *Thermotoga neapolitana*. *Int. J. Hydrogen Energy* **2010**, *35*, 13206–13213. [\[CrossRef\]](#)
50. Mars, A.E.; Veuskens, T.; Budde, M.A.W.; Van Doeveren, P.F.; Lips, S.J.; Bakker, R.R.; De Vrije, T.; Claassen, P.A.M. Biohydrogen Production from Untreated and Hydrolyzed Potato Steam Peels by the Extreme Thermophiles *Caldicellulosiruptor saccharolyticus* and *Thermotoga neapolitana*. *Int. J. Hydrogen Energy* **2010**, *35*, 7730–7737. [\[CrossRef\]](#)
51. Cappelletti, M.; Bucchi, G.; De Sousa Mendes, J.; Alberini, A.; Fedi, S.; Bertin, L.; Frascari, D. Biohydrogen Production from Glucose, Molasses and Cheese Whey by Suspended and Attached Cells of Four Hyperthermophilic Thermotoga Strains. *J. Chem. Technol. Biotechnol.* **2012**, *87*, 1291–1301. [\[CrossRef\]](#)
52. Saidi, R.; Liebgott, P.P.; Gannoun, H.; Ben Gaida, L.; Miladi, B.; Hamdi, M.; Bouallagui, H.; Auria, R. Biohydrogen Production from Hyperthermophilic Anaerobic Digestion of Fruit and Vegetable Wastes in Seawater: Simplification of the Culture Medium of *Thermotoga maritima*. *Waste Manag.* **2018**, *71*, 474–484. [\[CrossRef\]](#)
53. Saidi, R.; Liebgott, P.P.; Hamdi, M.; Auria, R.; Bouallagui, H. Enhancement of Fermentative Hydrogen Production by *Thermotoga maritima* through Hyperthermophilic Anaerobic Co-Digestion of Fruit-Vegetable and Fish Wastes. *Int. J. Hydrogen Energy* **2018**, *43*, 23168–23177. [\[CrossRef\]](#)
54. Saidi, R.; Hamdi, M.; Bouallagui, H. Hyperthermophilic Hydrogen Production in a Simplified Reaction Medium Containing Onion Wastes as a Source of Carbon and Sulfur. *Environ. Sci. Pollut. Res.* **2020**, *27*, 17382–17392. [\[CrossRef\]](#)

55. Amend, J.P.; Shock, E.L. Energetics of Overall Metabolic Reactions of Thermophilic and Hyperthermophilic Archaea and Bacteria. *FEMS Microbiol. Rev.* **2001**, *25*, 175–243. [\[CrossRef\]](#) [\[PubMed\]](#)
56. Pradhan, N.; Dipasquale, L.; d'Ippolito, G.; Panico, A.; Lens, P.N.L.; Esposito, G.; Fontana, A. Hydrogen Production by the Thermophilic Bacterium *Thermotoga neapolitana*. *Int. J. Mol. Sci.* **2015**, *16*, 12578–12600. [\[CrossRef\]](#) [\[PubMed\]](#)
57. Levin, D.B.; Pitt, L.; Love, M. Biohydrogen Production: Prospects and Limitations to Practical Application. *Int. J. Hydrogen Energy* **2004**, *29*, 173–185. [\[CrossRef\]](#)
58. Schut, G.J.; Adams, M.W.W. The Iron-Hydrogenase of *Thermotoga maritima* Utilizes Ferredoxin and NADH Synergistically: A New Perspective on Anaerobic Hydrogen Production. *J. Bacteriol.* **2009**, *191*, 4451–4457. [\[CrossRef\]](#)
59. Yu, J.; Varga, M.; Mityas, C.; Noll, K.M. Liposome-Mediated DNA Uptake and Transient Expression in *Thermotoga*. *Extremophiles* **2001**, *5*, 53–60. [\[CrossRef\]](#)
60. Xu, H.; Han, D.; Xu, Z. Expression of Heterologous Cellulases in *Thermotoga* Sp. Strain RQ2. *BioMed Res. Int.* **2015**, *2015*, 304523. [\[CrossRef\]](#)
61. Han, D.; Xu, Z. Development of a PyrE—Based Selective System for *Thermotoga* sp. *Extremophiles* **2017**, *21*, 297–306. [\[CrossRef\]](#)
62. Pradhan, N.; Dipasquale, L.; Panico, A.; Lens, P.N.L.; Esposito, G.; d'Ippolito, G.; Fontana, A. Hydrogen and Lactic Acid Synthesis by the Wild-Type and a Laboratory Strain of the Hyperthermophilic Bacterium *Thermotoga neapolitana* DSMZ 4359 under Capnophilic Lactic Fermentation Conditions. *Int. J. Hydrogen Energy* **2017**, *42*, 16023–16030. [\[CrossRef\]](#)
63. Grogan, D.W.; Carver, G.T.; Drake, J.W. Genetic fidelity under harsh conditions: Analysis of spontaneous mutation in the thermoacidophilic archaeon *Sulfolobus acidocaldarius*. *Proc. Natl. Acad. Sci. USA* **2001**, *98*, 7928–7933. [\[CrossRef\]](#)
64. Nguyen, T.A.D.; Pyo Kim, J.; Sun Kim, M.; Kwan Oh, Y.; Sim, S.J. Optimization of Hydrogen Production by Hyperthermophilic Eubacteria, *Thermotoga maritima* and *Thermotoga neapolitana* in Batch Fermentation. *Int. J. Hydrogen Energy* **2008**, *33*, 1483–1488. [\[CrossRef\]](#)
65. Munro, S.A.; Zinder, S.H.; Walker, L.P. The Fermentation Stoichiometry of *Thermotoga neapolitana* and Influence of Temperature, Oxygen, and PH on Hydrogen Production. *Biotechnol. Prog.* **2009**, *25*, 1035–1042. [\[CrossRef\]](#) [\[PubMed\]](#)
66. Cappelletti, M.; Zannoni, D.; Postec, A.; Ollivier, B. Members of the Order Thermotogales: From Microbiology to Hydrogen Production. In *Microbial BioEnergy: Hydrogen Production, Advances in Photosynthesis and Respiration*; Springer: Dordrecht, The Netherlands, 2014; Volume 38, pp. 197–224.
67. Pradhan, N.; d'Ippolito, G.; Dipasquale, L.; Esposito, G.; Panico, A.; Lens, P.N.L.; Fontana, A. Simultaneous Synthesis of Lactic Acid and Hydrogen from Sugars via Capnophilic Lactic Fermentation by *Thermotoga neapolitana* Cf *Capnolactica*. *Biomass Bioenergy* **2019**, *125*, 17–22. [\[CrossRef\]](#)
68. Dreschke, G.; d'Ippolito, G.; Panico, A.; Lens, P.N.L.; Esposito, G.; Fontana, A. Enhancement of Hydrogen Production Rate by High Biomass Concentrations of *Thermotoga Neapolitana*. *Int. J. Hydrogen Energy* **2018**, *43*, 13072–13080. [\[CrossRef\]](#)
69. Dreschke, G.; Papirio, S.; Panico, A.; Lens, P.N.L.; Esposito, G.; d'Ippolito, G.; Fontana, A. H₂-Rich Biogas Recirculation Prevents Hydrogen Supersaturation and Enhances Hydrogen Production by *Thermotoga neapolitana* Cf. *Capnolactica*. *Int. J. Hydrogen Energy* **2019**, *44*, 19698–19708. [\[CrossRef\]](#)
70. Dipasquale, L.; Pradhan, N.; d'Ippolito, G.; Fontana, A. Potential of Hydrogen Fermentative Pathways in Marine Thermophilic Bacteria: Dark Fermentation and Capnophilic Lactic Fermentation in *Thermotoga* and *Pseudothermotoga* Species. In *Grand Challenges in Marine Biotechnology*; Springer: Dordrecht, The Netherlands, 2018; pp. 217–235.
71. Nuzzo, G.; Landi, S.; Esercizio, N.; Manzo, E.; Fontana, A.; d'Ippolito, G. Capnophilic Lactic Fermentation from *Thermotoga neapolitana*: A Resourceful Pathway to Obtain Almost Enantiopure L-Lactic Acid. *Fermentation* **2019**, *5*, 34. [\[CrossRef\]](#)
72. Takahata, Y.; Nishijima, M.; Hoaki, T.; Maruyama, T. *Thermotoga petrophila* sp. Nov. and *Thermotoga naphthophila* sp. Nov., Two Hyperthermophilic Bacteria from the Kubiki Oil Reservoir in Niigata, Japan. *Int. J. Syst. Evol. Microbiol.* **2001**, *51*, 1901–1909. [\[CrossRef\]](#) [\[PubMed\]](#)
73. Mori, K.; Yamazoe, A.; Hosoyama, A.; Ohji, S.; Fujita, N.; Ishibashi, J.I.; Kimura, H.; Suzuki, K.I. *Thermotoga profunda* sp. Nov. and *Thermotoga caldifontis* sp. Nov., Anaerobic Thermophilic Bacteria Isolated from Terrestrial Hot Springs. *Int. J. Syst. Evol. Microbiol.* **2014**, *64*, 2128–2136. [\[CrossRef\]](#) [\[PubMed\]](#)
74. Jannasch, H.W.; Huber, R.; Belkin, S.; Stetter, K.O. *Thermotoga neapolitana* sp. Nov. of the Extremely Thermophilic, Eubacterial Genus *Thermotoga*. *Arch. Microbiol.* **1988**, *150*, 103–104. [\[CrossRef\]](#)
75. Balk, M.; Weijma, J.; Stams, A.J.M. *Thermotoga Lettingae* sp. Nov., a Novel Thermophilic, Methanol-Degrading Bacterium Isolated from a Thermophilic Anaerobic Reactor. *Int. J. Syst. Evol. Microbiol.* **2002**, *52*, 1361–1368. [\[CrossRef\]](#) [\[PubMed\]](#)
76. Ravot, G.; Magot, M.; Fardeau, M.L.; Patel, B.K.C.; Prensier, G.; Egan, A.; Garcia, J.L.; Ollivier, B. *Thermotoga elfii* sp. Nov., a Novel Thermophilic Bacterium from an African Oil-Producing Well. *Int. J. Syst. Bacteriol.* **1995**, *45*, 308–314. [\[CrossRef\]](#)
77. Fardeau, M.L.; Ollivier, B.C.; Patel, B.K.; Magot, M.; Thomas, P.; Rimbault, A.; Rocchiccioli, F.; Garcia, J.L. *Thermotoga hypogea* sp. Nov., a Xylanolytic, Thermophilic Bacterium from an Oil-Producing Well. *Int. J. Syst. Bacteriol.* **1997**, *147*, 51–56. [\[CrossRef\]](#) [\[PubMed\]](#)
78. Jeanthon, C.; Reysenbach, A.L.; L'Haridon, S.; Gambacorta, A.; Pace, N.R.; Glénat, P.; Prieur, D. *Thermotoga subterranea* sp. Nov., a New Thermophilic Bacterium Isolated from a Continental Oil Reservoir. *Arch. Microbiol.* **1995**, *164*, 91–97. [\[CrossRef\]](#) [\[PubMed\]](#)
79. Friedrich, A.B.; Antranikian, G. Keratin Degradation by *Fervidobacterium pennavorans*, a Novel Thermophilic Anaerobic Species of the Order Thermotogales. *Appl. Environ. Microbiol.* **1996**, *62*, 2875–2882. [\[CrossRef\]](#) [\[PubMed\]](#)

80. Huber, R.; Woese, C.R.; Langworthy, T.A.; Kristjansson, J.K.; Stetter, K.O. *Fervidobacterium islandicum* Sp. Nov., a New Extremely Thermophilic Eubacterium Belonging to the “Thermotogales”. *Arch. Microbiol.* **1990**, *154*, 105–111. [\[CrossRef\]](#)
81. Podosokorskaya, O.A.; Merkel, Y.A.; Kolganova, T.V.; Chernyh, N.A.; Miroshnichenko, M.L.; Bonch-Osmolovskaya, E.A.; Kublanov, I.V. *Fervidobacterium riparium* Sp. Nov., a Thermophilic Anaerobic Cellulolytic Bacterium Isolated from a Hot Spring. *Int. J. Syst. Evol. Microbiol.* **2011**, *61*, 2697–2701. [\[CrossRef\]](#)
82. Andrews, K.T.; Patel, B.K.C. *Fervidobacterium gondwanense* Sp. Nov., a New Thermophilic Anaerobic Bacterium Isolated from Nonvolcanically Heated Geothermal Waters of the Great Artesian Basin of Australia. *Int. J. Syst. Bacteriol.* **1996**, *46*, 265–269. [\[CrossRef\]](#)
83. Kanoksilapatham, W.; Pasomsup, P.; Keawram, P.; Cuecas, A.; Portillo, M.C.; Gonzalez, J.M. *Fervidobacterium thailandense* Sp. Nov., an Extremely Thermophilic Bacterium Isolated from a Hot Spring. *Int. J. Syst. Evol. Microbiol.* **2016**, *66*, 5023–5027. [\[CrossRef\]](#)
84. Cai, J.; Wang, Y.; Liu, D.; Zeng, Y.; Xue, Y.; Ma, Y.; Feng, Y. *Fervidobacterium changbaicum* Sp. Nov., a Novel Thermophilic Anaerobic Bacterium Isolated from a Hot Spring of the Changbai Mountains, China. *Int. J. Syst. Evol. Microbiol.* **2007**, *57*, 2333–2336. [\[CrossRef\]](#)
85. Huber, R.; Woese, C.R.; Langworthy, T.A.; Fricke, H.; Stetter, K.O. *Thermosipho africanus* Gen. Nov., Represents a New Genus of Thermophilic Eubacteria within the “Thermotogales”. *Syst. Appl. Microbiol.* **1989**, *12*, 32–37. [\[CrossRef\]](#)
86. Takai, K.; Horikoshi, K. *Thermosipho japonicus* Sp. Nov., an Extremely Thermophilic Bacterium Isolated from a Deep-Sea Hydrothermal Vent in Japan. *Extremophiles* **2000**, *4*, 9–17. [\[CrossRef\]](#)
87. L’Haridon, S.; Miroshnichenko, M.L.; Hippe, H.; Fardeau, M.L.; Bonch-Osmolovskaya, E.; Stackebrandt, E.; Jeanthon, C. *Thermosipho geolei* Sp. Nov., a Thermophilic Bacterium Isolated from a Continental Petroleum Reservoir in Western Siberia. *Int. J. Syst. Evol. Microbiol.* **2001**, *51*, 1327–1334. [\[CrossRef\]](#) [\[PubMed\]](#)
88. Podosokorskaya, O.A.; Kublanov, I.V.; Reysenbach, A.L.; Kolganova, T.V.; Bonch-Osmolovskaya, E.A. *Thermosipho affectus* Sp. Nov., a Thermophilic, Anaerobic, Cellulolytic Bacterium Isolated from a Mid-Atlantic Ridge Hydrothermal Vent. *Int. J. Syst. Evol. Microbiol.* **2011**, *61*, 1160–1164. [\[CrossRef\]](#) [\[PubMed\]](#)
89. Kuwabara, T.; Kawasaki, A.; Uda, I.; Sugai, A. *Thermosipho globiformans* Sp. Nov., an Anaerobic Thermophilic Bacterium That Transforms into Multicellular Spheroids with a Defect in Peptidoglycan Formation. *Int. J. Syst. Evol. Microbiol.* **2011**, *61*, 1622–1627. [\[CrossRef\]](#) [\[PubMed\]](#)
90. Antoine, E.; Cilia, V.; Meunier, J.R.; Guezennec, J.; Lesongeur, F.; Barbier, G. *Thermosipho melanesiensis* Sp. Nov., a New Thermophilic Anaerobic Bacterium Belonging to the Order Thermotogales, Isolated from Deep-Sea Hydrothermal Vents in the Southwestern Pacific Ocean. *Int. J. Syst. Bacteriol.* **1997**, *47*, 1118–1123. [\[CrossRef\]](#) [\[PubMed\]](#)
91. Podosokorskaya, O.A.; Bonch-Osmolovskaya, E.A.; Godfroy, A.; Gavrillov, S.N.; Beskorovaynaya, D.A.; Sokolova, T.G.; Kolganova, T.V.; Toshchakov, S.V.; Kublanov, I.V. *Thermosipho activus* Sp. Nov., a Thermophilic, Anaerobic, Hydrolytic Bacterium Isolated from a Deep-Sea Sample. *Int. J. Syst. Evol. Microbiol.* **2014**, *64*, 3307–3313. [\[CrossRef\]](#) [\[PubMed\]](#)
92. Urios, L.; Cueff-Gauchard, V.; Pignet, P.; Postec, A.; Fardeau, M.L.; Ollivier, B.; Barbier, G. *Thermosipho atlanticus* Sp. Nov., a Novel Member of the Thermotogales Isolated from a Mid-Atlantic Ridge Hydrothermal Vent. *Int. J. Syst. Evol. Microbiol.* **2004**, *54*, 1953–1957. [\[CrossRef\]](#)
93. L’Haridon, S.; Miroshnichenko, M.L.; Hippe, H.; Fardeau, M.L.; Stackebrandt, E.; Jeanthon, C. *Petrotoga olearia* Sp. Nov. and *Petrotoga sibirica* Sp. Nov., Two Thermophilic Bacteria Isolated from a Continental Petroleum Reservoir in Western Siberia. *Int. J. Syst. Evol. Microbiol.* **2002**, *52*, 1715–1722. [\[CrossRef\]](#)
94. Lien, T.; Madsen, M.; Rainey, F.A.; Birkeland, N.K. *Petrotoga mobilis* Sp. Nov., from a North Sea Oil-Production Well. *Int. J. Syst. Bacteriol.* **1998**, *48*, 1007–1013. [\[CrossRef\]](#)
95. Miranda-Tello, E.; Fardeau, M.L.; Joulian, C.; Magot, M.; Thomas, P.; Tholozan, J.L.; Olivier, B. *Petrotoga halophila* Sp. Nov., a Thermophilic, Moderately Halophilic, Fermentative Bacterium Isolated from an Offshore Oil Well in Congo. *Int. J. Syst. Evol. Microbiol.* **2007**, *57*, 40–44. [\[CrossRef\]](#)
96. Miranda-Tello, E.; Fardeau, M.L.; Thomas, P.; Ramirez, F.; Casalot, L.; Cayol, J.L.; Garcia, J.L.; Ollivier, B. *Petrotoga mexicana* Sp. Nov., a Novel Thermophilic, Anaerobic and Xylanolytic Bacterium Isolated from an Oil-Producing Well in the Gulf of Mexico. *Int. J. Syst. Evol. Microbiol.* **2004**, *54*, 169–174. [\[CrossRef\]](#)
97. Purwasena, I.A.; Sugai, Y.; Sasaki, K. *Petrotoga japonica* Sp. Nov., a Thermophilic, Fermentative Bacterium Isolated from Yabase Oilfield in Japan. *Arch. Microbiol.* **2014**, *196*, 313–321. [\[CrossRef\]](#) [\[PubMed\]](#)
98. Alain, K.; Marteinsson, V.T.; Miroshnichenko, M.L.; Bonch-Osmolovskaya, E.A.; Prieur, D.; Birrien, J.-L. *Marinitoga piezophila* Sp. Nov., a Rod-Shaped, Thermo-Piezophilic Bacterium Isolated under High Hydrostatic Pressure from a Deep-Sea Hydrothermal Vent. *Int. J. Syst. Evol. Microbiol.* **2002**, *52*, 1331–1339. [\[CrossRef\]](#) [\[PubMed\]](#)
99. Postec, A.; Ciobanu, M.; Birrien, J.L.; Bienvenu, N.; Prieur, D.; Le Romancer, M. *Marinitoga litoralis* Sp. Nov., a Thermophilic, Heterotrophic Bacterium Isolated from a Coastal Thermal Spring on Île Saint-Paul, Southern Indian Ocean. *Int. J. Syst. Evol. Microbiol.* **2010**, *60*, 1778–1782. [\[CrossRef\]](#) [\[PubMed\]](#)
100. Nunoura, T.; Oida, H.; Miyazaki, M.; Suzuki, Y.; Takai, K.; Horikoshi, K. *Marinitoga okinawensis* Sp. Nov., a Novel Thermophilic and Anaerobic Heterotroph Isolated from a Deep-Sea Hydrothermal Field, Southern Okinawa Trough. *Int. J. Syst. Evol. Microbiol.* **2007**, *57*, 467–471. [\[CrossRef\]](#) [\[PubMed\]](#)

101. Postec, A.; Le Breton, C.; Fardeau, M.L.; Lesongeur, F.; Pignet, P.; Querellou, J.; Ollivier, B.; Godfroy, A. *Marinitoga hydrogenitolerans* Sp. Nov., a Novel Member of the Order Thermotogales Isolated from a Black Smoker Chimney on the Mid-Atlantic Ridge. *Int. J. Syst. Evol. Microbiol.* **2005**, *55*, 1217–1221. [[CrossRef](#)] [[PubMed](#)]
102. Steinsbu, B.O.; Røyseth, V.; Thorseth, I.H.; Steen, I.H. *Marinitoga arctica* Sp. Nov., a Thermophilic, Anaerobic Heterotroph Isolated from a Mid-Ocean Ridge Vent Field. *Int. J. Syst. Evol. Microbiol.* **2016**, *66*, 5070–5076. [[CrossRef](#)]
103. Nunoura, T.; Hirai, M.; Imachi, H.; Miyazaki, M.; Makita, H.; Hirayama, H.; Furushima, Y.; Yamamoto, H.; Takai, K. *Kosmotoga arenicorallina* Sp. Nov. a Thermophilic and Obligately Anaerobic Heterotroph Isolated from a Shallow Hydrothermal System Occurring within a Coral Reef, Southern Part of the Yaeyama Archipelago, Japan, Reclassification of *Thermococcoides shengliensis*. *Arch. Microbiol.* **2010**, *192*, 811–819. [[CrossRef](#)] [[PubMed](#)]
104. L'Haridon, S.; Jiang, L.; Alain, K.; Chalopin, M.; Rouxel, O.; Beauverger, M.; Xu, H.; Shao, Z.; Jebbar, M. *Kosmotoga pacifica* Sp. Nov., a Thermophilic Chemoorganoheterotrophic Bacterium Isolated from an East Pacific Hydrothermal Sediment. *Extremophiles* **2014**, *18*, 81–88. [[CrossRef](#)]
105. Van Ooteghem, S.A.; Beer, S.K.; Yue, P.C. Hydrogen Production by the Thermophilic Bacterium *Thermotoga neapolitana*. *Appl. Biochem. Biotechnol.* **2002**, *98–100*, 177–189. [[CrossRef](#)]
106. Eriksen, N.T.; Nielsen, T.M.; Iversen, N. Hydrogen Production in Anaerobic and Microaerobic *Thermotoga neapolitana*. *Biotechnol. Lett.* **2008**, *30*, 103–109. [[CrossRef](#)]
107. Boileau, C.; Auria, R.; Davidson, S.; Casalot, L.; Christen, P.; Liebgott, P.P.; Combet-Blanc, Y. Hydrogen Production by the Hyperthermophilic Bacterium *Thermotoga maritima* Part I: Effects of Sulfured Nutriments, with Thiosulfate as Model, on Hydrogen Production and Growth. *Biotechnol. Biofuels* **2016**, *9*, 1–17. [[CrossRef](#)] [[PubMed](#)]
108. Van Niel, E.W.J.; Budde, M.A.W.; De Haas, G.; van der Wal, F.J.; Claassen, P.A.M.; Stams, A.J.M. Distinctive Properties of High Hydrogen Producing Extreme Thermophiles, *Caldicellulosiruptor saccharolyticus* and *Thermotoga elfii*. *Int. J. Hydrogen Energy* **2002**, *27*, 1391–1398. [[CrossRef](#)]
109. Hawkes, F.; Hussy, I.; Kyazze, G.; Dinsdale, R.; Hawkes, D. Continuous Dark Fermentative Hydrogen Production by Mesophilic Microflora: Principles and Progress. *Int. J. Hydrogen Energy* **2007**, *32*, 172–184. [[CrossRef](#)]
110. Nguyen, T.A.D.; Han, S.J.; Kim, J.P.; Kim, M.S.; Sim, S.J. Hydrogen Production of the Hyperthermophilic Eubacterium, *Thermotoga neapolitana* under N₂ Sparging Condition. *Bioresour. Technol.* **2010**, *101*, S38–S41. [[CrossRef](#)]
111. Karadagli, F.; Marcus, A.K.; Rittmann, B.E. Role of Hydrogen (H₂) Mass Transfer in Microbiological H₂-Threshold Studies. *Biodegradation* **2019**, *30*, 113–125. [[CrossRef](#)]
112. Nath, K.; Das, D. Improvement of Fermentative Hydrogen Production: Various Approaches. *Appl. Microbiol. Biotechnol.* **2004**, *65*, 520–529. [[CrossRef](#)]
113. Ngo, T.A.; Mi-sun, K.; Sim, S.J. Thermophilic Hydrogen Fermentation Using *Thermotoga neapolitana* DSM 4359 by Fed-Batch Culture. *Int. J. Hydrogen Energy* **2011**, *36*, 14014–14023. [[CrossRef](#)]
114. Pradhan, N.; Dipasquale, L.; d'Ippolito, G.; Fontana, A.; Panico, A.; Lens, P.N.L.; Pirozzi, F.; Esposito, G. Kinetic modeling of fermentative hydrogen production by *Thermotoga neapolitana*. *Int. J. Hydrogen Energy* **2016**, *41*, 1–10. [[CrossRef](#)]
115. Mizuno, O.; Dinsdale, R.; Hawkes, F.R.; Hawkes, D.L. Enhancement of Hydrogen Production from Glucose by Nitrogen Gas Sparging. *Bioresour. Technol.* **2000**, *73*, 59–65. [[CrossRef](#)]
116. Ngo, T.A.; Kim, M.S.; Sim, S.J. High-Yield Biohydrogen Production from Biodiesel Manufacturing Waste by *Thermotoga neapolitana*. *Int. J. Hydrogen Energy* **2011**, *36*, 5836–5842. [[CrossRef](#)]
117. D'Ippolito, G.; Dipasquale, L.; Fontana, A. Recycling of Carbon Dioxide and Acetate as Lactic Acid by the Hydrogen-Producing Bacterium *Thermotoga neapolitana*. *ChemSuschem* **2014**, *7*, 2678–2683. [[CrossRef](#)] [[PubMed](#)]
118. Pradhan, N.; Dipasquale, L.; d'Ippolito, G.; Fontana, A.; Panico, A.; Pirozzi, F.; Lens, P.N.L.; Esposito, G. Model Development and Experimental Validation of Capnophilic Lactic Fermentation and Hydrogen Synthesis by *Thermotoga neapolitana*. *Water Res.* **2016**, *99*, 225–234. [[CrossRef](#)] [[PubMed](#)]
119. Dipasquale, L.; Adessi, A.; d'Ippolito, G.; Rossi, F.; Fontana, A.; De Philippis, R. Introducing Capnophilic Lactic Fermentation in a Combined Dark-Photo Fermentation Process: A Route to Unparalleled H₂ Yields. *Appl. Microbiol. Biotechnol.* **2015**, *99*, 1001–1010. [[CrossRef](#)] [[PubMed](#)]
120. D'Ippolito, G.; Landi, S.; Esercizio, N.; Lanzilli, M.; Vastano, M.; Dipasquale, L.; Pradhan, N.; Fontana, A. CO₂-Induced Transcriptional Reorganization: Molecular Basis of Capnophilic Lactic Fermentation in *Thermotoga neapolitana*. *Front. Microbiol.* **2020**, *11*, 171. [[CrossRef](#)] [[PubMed](#)]
121. Van Ooteghem, S.A.; Jones, A.; Van Der Lelie, D.; Dong, B.; Mahajan, D. H₂ Production and Carbon Utilization by *Thermotoga neapolitana* under Anaerobic and Microaerobic Growth Conditions. *Biotechnol. Lett.* **2004**, *26*, 1223–1232. [[CrossRef](#)] [[PubMed](#)]
122. Ngo, T.A.; Sim, S.J. Dark Fermentation of Hydrogen from Waste Glycerol Using Hyperthermophilic Eubacterium *Thermotoga neapolitana*. *Environ. Prog. Sustain. Energy* **2011**, *31*, 466–473. [[CrossRef](#)]
123. Juszczak, A.; Aono, S.; Adams, M.W.W. The Extremely Thermophilic Eubacterium, *Thermotoga maritima* Contains a Novel Iron-Hydrogenase Whose Cellular Activity Is Dependent Upon Tungsten. *J. Bacteriol. Chem.* **1991**, *226*, 13834–13841. [[PubMed](#)]
124. Rusch, A.; Walpersdorf, E.; DeBeer, D.; Gurrier, S.; Amend, P.J. Microbial Communities near the Oxidic/Anoxic Interface in the Hydrothermal System of Vulcano Island, Italy. *Chem. Geol.* **2005**, *224*, 169–182. [[CrossRef](#)]
125. Belkin, S.; Wirsén, C.O.; Jannasch, H.W. A New Sulfur-Reducing, Extremely Thermophilic Eubacterium from a Submarine Thermal Vent. *Appl. Environ. Microbiol.* **1986**, *51*, 1180–1185. [[CrossRef](#)]

126. Childers, S.E.; Vargas, M.; Noll, K.M. Improved Methods for Cultivation of the Extremely Thermophilic Bacterium *Thermotoga neapolitana*. *Appl. Environ. Microbiol.* **1992**, *58*, 3949–3953. [[CrossRef](#)]
127. Le Fourn, C.; Fardeau, M.L.; Ollivier, B.; Lojou, E.; Dolla, A. The Hyperthermophilic Anaerobe *Thermotoga maritima* Is Able to Cope with Limited Amount of Oxygen: Insights into Its Defence Strategies. *Environ. Microbiol.* **2008**, *10*, 1877–1887. [[CrossRef](#)] [[PubMed](#)]
128. Tosatto, S.C.E.; Toppo, S.; Carbonera, D.; Giacometti, G.M.; Costantini, P. Comparative Analysis of [FeFe] Hydrogenase from *Thermotogales* Indicates the Molecular Basis of Resistance to Oxygen Inactivation. *Int. J. Hydrogen Energy* **2008**, *33*, 570–578. [[CrossRef](#)]
129. Lakhal, R.; Auria, R.; Davidson, S.; Ollivier, B.; Dolla, A.; Hamdi, M.; Combet-Blanc, Y. Effect of Oxygen and Redox Potential on Glucose Fermentation in *Thermotoga maritima* under Controlled Physicochemical Conditions. *Int. J. Microbiol.* **2010**, *2010*, 896510. [[CrossRef](#)] [[PubMed](#)]
130. Kaslin, S.A.; Childers, S.E.; Noll, K.M. Membrane-Associated Redox Activities in *Thermotoga neapolitana*. *Arch. Microbiol.* **1998**, *170*, 297–303. [[CrossRef](#)] [[PubMed](#)]
131. Yang, X.; Ma, K. Purification and Characterization of an NADH Oxidase from Extremely Thermophilic Anaerobic Bacterium *Thermotoga hypogea*. *Arch. Microbiol.* **2005**, *183*, 331–337. [[CrossRef](#)] [[PubMed](#)]
132. Yang, X.; Ma, K. Characterization of an Exceedingly Active NADH Oxidase from the Anaerobic Hyperthermophilic Bacterium *Thermotoga maritima*. *J. Bacteriol.* **2007**, *189*, 3312–3317. [[CrossRef](#)]
133. Mangayil, R.; Aho, T.; Karp, M.; Santala, V. Improved Bioconversion of Crude Glycerol to Hydrogen by Statistical Optimization of Media Components. *Renew. Energy* **2015**, *75*, 583–589. [[CrossRef](#)]
134. Abdullah, M.F.; Md Jahim, J.; Abdul, P.M.; Mahmod, S.S. Effect of Carbon/Nitrogen Ratio and Ferric Ion on the Production of Biohydrogen from Palm Oil Mill Effluent (POME). *Biocatal. Agric. Biotechnol.* **2020**, *23*, 101445. [[CrossRef](#)]
135. Munro, S.A.; Choe, L.; Zinder, S.H.; Lee, K.H.; Walker, L.P. Proteomic and Physiological Experiments to Test *Thermotoga neapolitana* Constraint-Based Model Hypotheses of Carbon Source Utilization. *Biotechnol. Prog.* **2012**, *28*, 312–318. [[CrossRef](#)]
136. Rinker, K.D.; Kelly, R.M. Effect of Carbon and Nitrogen Sources on Growth Dynamics and Exopolysaccharide Production for the Hyperthermophilic Archaeon *Thermococcus litoralis* and Bacterium *Thermotoga maritima*. *Biotechnol. Bioeng.* **2000**, *69*, 537–547. [[CrossRef](#)]
137. Van Niel, E.W.J.; Hahn-Hägerdal, B. Nutrient Requirements of Lactococci in Defined Growth Media. *Appl. Microbiol. Biotechnol.* **1999**, *52*, 617–627. [[CrossRef](#)]
138. Maru, B.T.; Bielen, A.A.M.; Kengen, S.W.M.; Constantí, M.; Medinaa, F. Biohydrogen Production from Glycerol Using *Thermotoga* Spp. *Energy Procedia* **2012**, *29*, 300–307. [[CrossRef](#)]
139. Kengen, S.W.M.; Stams, A.J.M. Formation of L-Alanine as a Reduced End Product in Carbohydrate Fermentation by the Hyperthermophilic Archaeon *Pyrococcus furiosus*. *Arch. Microbiol.* **1994**, *161*, 168–175. [[CrossRef](#)]
140. Weiner, R.; Langille, S.; Quintero, E. Structure, Function, and Immu- Nochemistry of Bacterial Exopolysaccharides. *J. Ind. Microbiol.* **1995**, *15*, 339–346. [[CrossRef](#)] [[PubMed](#)]
141. Pierra, M.; Trably, E.; Godon, J.J.; Bernet, N. Fermentative Hydrogen Production under Moderate Halophilic Conditions. *Int. J. Hydrogen Energy* **2014**, *39*, 7508–7517. [[CrossRef](#)]
142. Liu, Q.; Chen, W.; Zhang, X.; Yu, L.; Zhou, J.; Xu, Y.; Qian, G. Phosphate Enhancing Fermentative Hydrogen Production from Substrate with Municipal Solid Waste Composting Leachate as a Nutrient. *Bioresour. Technol.* **2015**, *190*, 431–437. [[CrossRef](#)] [[PubMed](#)]
143. Chang, F.; Lin, C. Calcium Effect on Fermentative Hydrogen Production in an Anaerobic Up-Flow Sludge Blanket System. *Water Sci. Technol.* **2006**, *54*, 105–112. [[CrossRef](#)]
144. Ravot, G.; Ollivier, B.; Magot, M.; Patel, B.K.C.; Crolet, J.L.; Fardeau, M.L.; Garcia, J.L. Thiosulfate Reduction, an Important Physiological Feature Shared by Members of the Order Thermotogales. *Appl. Environ. Microbiol.* **1995**, *61*, 2053–2055. [[CrossRef](#)]
145. Ravot, G.; Ollivier, B.; Fardeau, M.L.; Patel, B.K.C.; Andrews, K.T.; Magot, M.; Garcia, J.L. L-Alanine Production from Glucose Fermentation by Hyperthermophilic Members of the Domains Bacteria and Archaea: A Remnant of an Ancestral Metabolism? *Appl. Environ. Microbiol.* **1996**, *62*, 2657–2659. [[CrossRef](#)]
146. Ainala, S.K.; Seol, E.; Kim, J.R.; Park, S. Effect of Culture Medium on Fermentative and CO-Dependent H₂ Production Activity in *Citrobacter amalonaticus* Y19. *Int. J. Hydrogen Energy* **2016**, *41*, 6734–6742. [[CrossRef](#)]
147. Huber, R.; Stetter, K.O. The Order Thermotogales. In *The Prokaryotes*; Springer: New York, NY, USA, 1992; pp. 93809–93815.
148. Hania, W.B.; Fadhlou, K.; Brochier-Armanet, C.; Persillon, C.; Postec, A.; Hamdi, M.; Dolla, A.; Ollivier, B.; Fardeau, M.L.; Le Mer, J.; et al. Draft Genome Sequence of Mesotoga Strain PhosAC3, a Mesophilic Member of the Bacterial Order Thermotogales, Isolated from a Digestor Treating Phosphogypsum in Tunisia. *Stand. Genom. Sci.* **2015**, *10*, 1–7. [[CrossRef](#)] [[PubMed](#)]
149. Nesbø, C.L.; Charchuk, R.; Pollo, S.M.J.; Budwill, K.; Kublanov, I.V.; Haverkamp, T.H.A.; Foght, J. Genomic Analysis of the Mesophilic Thermotogae Genus *Mesotoga* Reveals Phylogeographic Structure and Genomic Determinants of Its Distinct Metabolism. *Environ. Microbiol.* **2019**, *21*, 456–470. [[CrossRef](#)] [[PubMed](#)]
150. Tao, Z.; Yang, Q.; Yao, F.; Huang, X.; Wu, Y.; Du, M.; Chen, S.; Liu, X.; Li, X.; Wang, D. The Inhibitory Effect of Thiosulfate on Volatile Fatty Acid and Hydrogen Production from Anaerobic Co-Fermentation of Food Waste and Waste Activated Sludge. *Bioresour. Technol.* **2020**, *297*, 122428. [[CrossRef](#)] [[PubMed](#)]
151. Slobodkin, A.I. Thermophilic Microbial Metal Reduction. *Microbiologia* **2005**, *74*, 581–595. [[CrossRef](#)]

-
152. Gomez-Romero, J.; Gonzalez-Garcia, A.; Chairez, I.; Torres, L.; García-Peña, E.I. Selective Adaptation of an Anaerobic Microbial Community: Biohydrogen Production by Co-Digestion of Cheese Whey and Vegetables Fruit Waste. *Int. J. Hydrogen Energy* **2014**, *39*, 12541–12550. [[CrossRef](#)]
 153. Llanos, J.; Capasso, C.; Parisi, E.; Prieur, D.; Jeanthon, C. Susceptibility to Heavy Metals and Cadmium Accumulation in Aerobic and Anaerobic Thermophilic Microorganisms Isolated from Deep-Sea Hydrothermal Vents. *Curr. Microbiol.* **2000**, *41*, 0201–0205. [[CrossRef](#)]
 154. Vargas, M.; Kashefi, K.; Blunt-harris, E.L.; Lovley, D.R. Fe (III) Reduction on Early Earth. *Nature* **1998**, *395*, 65–67. [[CrossRef](#)]
 155. Balch, W.E.; Fox, G.E.; Magrum, L.J.; Woese, C.R.; Wolfe, R.S. Methanogens: Reevaluation of a Unique Biological Group. *Microbiol. Rev.* **1979**, *43*, 260–296. [[CrossRef](#)]
 156. Lee, Y.J.; Miyahara, T.; Noike, T. Effect of Iron Concentration on Hydrogen Fermentation. *Bioresour. Technol.* **2001**, *80*, 227–231. [[CrossRef](#)]
 157. Zhang, Y.; Shen, J. Effect of Temperature and Iron Concentration on the Growth and Hydrogen Production of Mixed Bacteria. *Int. J. Hydrogen Energy* **2006**, *31*, 441–446. [[CrossRef](#)]
 158. Ying, X.; Wang, Y.; Badiei, H.R.; Karanassios, V.; Ma, K. Purification and Characterization of an Iron-Containing Alcohol Dehydrogenase in Extremely Thermophilic Bacterium *Thermotoga hypogea*. *Arch. Microbiol.* **2007**, *187*, 499–510. [[CrossRef](#)] [[PubMed](#)]
 159. Trivedi, S.; Rao, S.R.; Gehlot, H.S. Nucleic Acid Stability in Thermophilic Prokaryotes: A Review Nucleic Acid Stability in Thermophilic Prokaryotes: A Review. *J. Cell Mol. Biol.* **2005**, *4*, 61–69.